

Periodic Delta Function, and Expansion in Chebyshev Polynomials

H. Vic Dannon
vic0@comcast.net
July, 2012

Abstract Let $f(x)$ be defined on $[-1,1]$, so that $f(1) = f(-1)$.
 $T_n(x) = \cos(n\theta_x)$, $\theta_x = \arccos x + 2\pi m$, or $\theta_x = -\arccos x + 2\pi m$
where m is an integer, are the Chebyshev Polynomials on $[-1,1]$.

$$T_0(x) = 1, T_1(x) = x, T_2(x) = 2x^2 - 1, T_3(x) = 4x^3 - 3x, \dots$$

The Chebyshev Series associated with $f(x)$ is

$$\frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + a_2 T_2(x) + \dots$$

where

$$a_n = \frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_n(\xi) d\xi$$

are the Chebyshev Series coefficients.

The Chebyshev Series Theorem supplies the conditions under which the Chebyshev Series associated with $f(x)$ equals $f(x)$.

It is believed to hold in the Calculus of Limits for smooth enough function. In fact,

*The Theorem cannot be proved in the Calculus of Limits
under any conditions,*

because the summation of the Chebyshev Series requires integration over the periodically singular Chebyshev Kernel.

Plots of partial sums of the Chebyshev Series speak volumes about the sensibility of the claims to have infinities bound by epsilons.

In Infinitesimal Calculus, the Chebyshev Kernel

$$\frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + .. + T_n(\xi)T_n(x) + ... \right\}$$

is the Periodic Delta Function,

$$\delta_{Periodic}(\theta_\xi - \theta_x) = ... + \delta(\theta_\xi - \theta_x + 2\pi) + \delta(\theta_\xi - \theta_x) + \delta(\theta_\xi - \theta_x - 2\pi) + ...$$

It equals its Chebyshev Series, and the Chebyshev Series associated with any hyper-real integrable $f(x)$, equals $f(x)$

Keywords: Infinitesimal, Infinite-Hyper-Real, Hyper-Real, infinite Hyper-real, Infinitesimal Calculus, Delta Function, Chebyshev Polynomials, Chebyshev Coefficients, Periodic Delta Function, Delta Comb, Chebyshev Series, Chebyshev Kernel, Expansion in Chebyshev Polynomials,

2000 Mathematics Subject Classification 26E35; 26E30;
26E15; 26E20; 26A06; 26A12; 03E10; 03E55; 03E17; 03H15;
46S20; 97I40; 97I30.

Contents

0. The Origin of the Chebyshev Series Theorem
1. Divergence of the Chebyshev Kernel in the Calculus of Limits
2. Hyper-real line.
3. Integral of a Hyper-real Function
4. Delta Function
5. Periodic Delta Function, $\delta_{Periodic}(\theta_\xi - \theta_x)$
6. Convergent Series
7. Chebyshev Sequence and $\delta_{Periodic}(\theta_\xi - \theta_x)$
8. Chebyshev Kernel and $\delta_{Periodic}(\theta_\xi - \theta_x)$.
9. Chebyshev Series of $\delta_{Periodic}(\theta_\xi - \theta_x)$
10. Chebyshev Series Theorem

References

The Origin of the Chebyshev Series

Theorem

The Chebyshev Polynomials $T_n(x) = \cos(n\theta_x)$, $\theta_x = \arccos x$, on $[-1,1]$, $T_0(x) = 1$, $T_1(x) = x$, $T_2(x) = 2x^2 - 1$, $T_3(x) = 4x^3 - 3x, \dots$ are the orthogonal functions generated in the Gram-Schmidt orthogonalization so that,

$$\frac{1}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_0(\theta_x) T_0(\theta_x) dx = 1,$$

$$\frac{2}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_m(\theta_x) T_n(\theta_x) dx = \delta_{mn}, \quad m, n = 1, 2, 3, \dots$$

The Chebyshev Polynomials can be generated by expanding

$$\frac{1 - x\alpha}{1 - 2x\alpha + \alpha^2} = T_0(x) + T_1(x)\alpha + T_2(x)\alpha^2 + T_3(x)\alpha^3 + \dots$$

$$\frac{1 - x\alpha}{1 - \underbrace{(2x - \alpha)\alpha}_q} = (1 - x\alpha)[1 + q + q^2 + \dots],$$

$$= (1 - x\alpha) \left[1 + (2x - \alpha)\alpha + (2x - \alpha)^2 \alpha^2 + (2x - \alpha)^3 \alpha^3 + \dots \right]$$

$$= 1 + (2x - \alpha)\alpha + (2x - \alpha)^2 \alpha^2 + (2x - \alpha)^3 \alpha^3 + \dots$$

$$-x\alpha - x(2x - \alpha)\alpha^2 - x(2x - \alpha)^2 \alpha^3 - x(2x - \alpha)^3 \alpha^4 + \dots$$

the coefficient of α^0 is

$$T_0(x) = 1,$$

the coefficient of α^1 is

$$T_1(x) = x,$$

the coefficient of α^2 is

$$T_2(x) = 2x^2 - 1,$$

the coefficient of α^3 is

$$T_3(x) = 4x^3 - 3x,$$

.....

0.1 Chebyshev

established that

$$\max_{-1 \leq x \leq 1} |T_n(x)| \leq \max_{-1 \leq x \leq 1} 2^{n-1} |x^n + a_{n-1}x^{n-1} + \dots + a_0|$$

That is, the maximal value of $|T_n(x)|$ on $[-1,1]$ is smaller than the maximal value of any $2^{n-1} |x^n + a_{n-1}x^{n-1} + \dots + a_0|$ on $[-1,1]$, [Krylov].

0.2 Chebyshev Differential Equation

$$(1 - x^2)y''(x) - xy'(x) + n^2y(x) = 0,$$

for $-1 \leq x \leq 1$, and $n = 0,1,2,3,\dots$

Substituting in it

$$y(x) = c_0 + c_1x + c_2x^2 + \dots + c_lx^l + c_{l+1}x^{l+1} + c_{l+2}x^{l+2} + \dots,$$

we have

$$\underbrace{D_x^2 \sum_{l=0}^{l=\infty} c_l x^l}_{\sum_{l=2}^{l=\infty} (l-1)lc_l x^{l-2}} - x^2 \underbrace{D_x^2 \sum_{l=0}^{l=\infty} c_l x^l}_{\sum_{l=2}^{l=\infty} (l-1)lc_l x^{l-2}} - x \underbrace{D_x \sum_{l=0}^{l=\infty} c_l x^l}_{\sum_{l=1}^{l=\infty} lc_l x^{l-1}} + n^2 \sum_{l=0}^{l=\infty} c_l x^l = 0,$$

$$\sum_{l=0}^{l=\infty} \{(l+1)(l+2)c_{l+2} - l(l-1)c_l - lc_l + n^2 c_l\} x^l = 0,$$

$$(l+1)(l+2)c_{l+2} - l(l-1)c_l - lc_l + n^2 c_l = 0,$$

$$c_{l+2} = \frac{l(l-1) + l - n^2}{(l+1)(l+2)} c_l$$

$$c_{l+2} = (-1) \frac{(n-l)(n+l)}{(l+1)(l+2)} c_l$$

The solution is

$$\begin{aligned} y(x) &= c_0 + c_1 x - c_0 \frac{n \cdot n}{1 \cdot 2} x^2 - c_1 \frac{(n-1)(n+1)}{1 \cdot 2 \cdot 3} x^3 + \\ &\quad + c_0 \frac{n \cdot n(n-2)(n+2)}{1 \cdot 2 \cdot 3 \cdot 4} x^4 + c_1 \frac{(n-1)(n+1)(n-3)(n+3)}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} x^5 + \dots \\ &= c_0 \left\{ 1 - \frac{n \cdot n}{1 \cdot 2} x^2 + \frac{n \cdot n(n-2)(n+2)}{1 \cdot 2 \cdot 3 \cdot 4} x^4 - \dots \right\} + \\ &\quad + c_1 x \left\{ 1 - \frac{(n-1)(n+1)}{1 \cdot 2 \cdot 3} x^2 + \frac{(n-1)(n+1)(n-3)(n+3)}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} x^4 + \dots \right\}. \end{aligned}$$

For $n = 2k$, the c_0 series terms vanish for

$$2k + 2, 2k + 4, \dots,$$

and we obtain the $T_{2k}(x)$ Chebyshev Polynomials.

for $n = 2k + 1$, the c_1 series terms vanish for

$$2k + 3, 2k + 5, \dots$$

and we obtain the $P_{2k+1}(x)$ Chebyshev Polynomials.

0.3 The Chebyshev Series Associated with a periodic $f(x)$

Let $f(x)$ be integrable on $[-1,1]$, so that $f(1) = f(-1)$, and let

$T_n(x) = \cos(n\theta_x)$, $\theta_x = \arccos x$, be the Chebyshev Polynomials on $[-1,1]$, $T_0(x) = 1$, $T_1(x) = x$, $T_2(x) = 2x^2 - 1$, $T_3(x) = 4x^3 - 3x, \dots$

The Polynomials are orthogonal on $[-1,1]$. That is,

$$\frac{1}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_0(\theta_x) T_0(\theta_x) dx = 1,$$

$$\frac{2}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_m(\theta_x) T_n(\theta_x) dx = \delta_{mn}, \quad m, n = 1, 2, 3, \dots$$

If $f(x)$ can be expanded in the Chebyshev Polynomials,

$$f(x) = \frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + a_2 T_2(x) + \dots,$$

Then,

$$\begin{aligned} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} f(x) T_n(x) dx &= \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} \left\{ \frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + a_2 T_2(x) + \dots \right\} T_n(x) dx \\ &= a_0 \underbrace{\frac{1}{2} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_0(x) T_n(x) dx}_{\frac{\pi}{2} \delta_{0n}} + a_1 \underbrace{\int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_1(x) T_n(x) dx}_{\frac{\pi}{2} \delta_{1n}} + \dots \end{aligned}$$

$$\begin{aligned}
& + a_2 \underbrace{\int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} T_2(x) T_n(x) dx}_{\frac{\pi}{2} \delta_{2n}} + \dots \\
& = a_n \frac{\pi}{2}.
\end{aligned}$$

Thus, the Chebyshev coefficients are

$$a_n = \frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_n(\xi) d\xi.$$

The Chebyshev Series associated with $f(x)$ is

$$\frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + a_2 T_2(x) + \dots$$

For $-1 \leq x \leq 1$,

$$x = \cos \theta_x,$$

where

$$\theta_x = \arccos x + 2\pi m,$$

or

$$\theta_x = -\arccos x + 2\pi m.$$

and m is an integer

Thus, the Chebyshev Polynomials,

$$T_n(x) = \cos(n\theta_x),$$

are periodic in θ , with period 2π .

Therefore, the Chebyshev Series is periodic in θ , with period 2π .

The Chebyshev Series Theorem supplies the conditions under which the Chebyshev Series associated with $f(x)$ equals $f(x)$.

Then, *the function must be periodic too.*

1.

Divergence of the Chebyshev Kernel in the Calculus of Limits

Calculus of Limits Conditions for the Chebyshev Series to equal its function reflect the belief that a smooth enough function equals its Chebyshev Series.

In fact, in the Calculus of Limits, no smoothness of the function guarantees even the convergence of the Chebyshev Series.

In the Calculus of Limits, the Chebyshev Series is the limit of the sequence of Partial Sums

$$\begin{aligned}
 \mathcal{C}_{chebyshev} \mathcal{S}_n \{f(x)\} &= \frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + \dots + a_n T_n(x) \\
 &= \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} \frac{1}{2} f(\xi) T_0(\xi) d\xi \right) P_0(x) + \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_1(\xi) d\xi \right) T_1(x) + \dots \\
 &\quad \dots + \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_n(\xi) d\xi \right) T_n(x) \\
 &= \int_{\xi=-1}^{\xi=1} f(\xi) \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) \dots + T_n(\xi) T_n(x) \right\} d\xi.
 \end{aligned}$$

As $n \rightarrow \infty$, the Chebyshev Sequence

$$\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi)T_0(x) + T_1(\xi)T_1(x) \dots + T_n(\xi)T_n(x) \right\}$$

becomes the Chebyshev Kernel,

$$\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi)T_0(x) + T_1(\xi)T_1(x) \dots + T_n(\xi)T_n(x) + \dots \right\},$$

Substituting $T_n(x) = \cos(n\theta_x)$,

$$\begin{aligned} & \frac{1}{2} T_0(\xi)T_0(x) + T_1(\xi)T_1(x) + T_2(\xi)T_2(x) + \dots = \\ & = \frac{1}{2} + \cos \theta_\xi \cos \theta_x + \cos 2\theta_\xi \cos 2\theta_x + \dots \\ & = \frac{1}{2} + \frac{1}{2} \left\{ \cos(\theta_\xi + \theta_x) + \cos(\theta_\xi - \theta_x) \right\} + \\ & \quad + \frac{1}{2} \left\{ \cos 2(\theta_\xi + \theta_x) + \cos 2(\theta_\xi - \theta_x) \right\} + \dots \\ & = \frac{1}{2} \left\{ \frac{1}{2} + \cos(\theta_\xi + \theta_x) + \cos 2(\theta_\xi + \theta_x) + \dots \right\} + \\ & \quad + \frac{1}{2} \left\{ \frac{1}{2} + \cos(\theta_\xi - \theta_x) + \cos 2(\theta_\xi - \theta_x) + \dots \right\}. \end{aligned}$$

To show that each series vanishes for $\xi \neq x$, we follow an Euler type argument [Hardy, p.2]. We have

$$\begin{aligned} s &= 1 + e^{i\theta} + e^{i2\theta} + e^{i3\theta} + \dots \\ &= 1 + e^{i\theta} \underbrace{\left[1 + e^{i\theta} + e^{i2\theta} + \dots \right]}_s \end{aligned}$$

For any $\theta \neq 2\pi m$,

$$e^{i\theta} \neq 1.$$

Thus, $s \neq 1 + 1 + \dots$, and

$$s = \frac{1}{1 - e^{i\theta}}$$

$$\begin{aligned}
&= \frac{e^{-\frac{1}{2}i\theta}}{e^{-\frac{1}{2}i\theta} - e^{\frac{1}{2}i\theta}} \\
&= \frac{\cos \frac{1}{2}\theta - i \sin \frac{1}{2}\theta}{-2i \sin \frac{1}{2}\theta} \\
&= i \frac{1}{2} \cot \frac{1}{2}(\theta_\xi - \theta_x) + \frac{1}{2}
\end{aligned}$$

Therefore,

$$\begin{aligned}
\operatorname{Re}[s] &= \frac{1}{2}, \\
\operatorname{Re}\left(1 + e^{i\theta} + e^{i2\theta} + e^{i3\theta} + \dots\right) &= \frac{1}{2}, \\
1 + \cos \theta + \cos 2\theta + \cos 3\theta + \dots &= \frac{1}{2}.
\end{aligned}$$

That is, for any $\theta \neq 2\pi m$,

$$\frac{1}{2} + \cos \theta + \cos 2\theta + \cos 3\theta + \dots = 0.$$

For $\xi \neq x$,

$$\begin{aligned}
\underbrace{\arccos \xi}_{\theta_\xi} \neq \underbrace{\arccos x}_{\theta_x} + 2\pi m &\Rightarrow \theta_\xi - \theta_x \neq 2\pi m \Rightarrow \\
&\Rightarrow \frac{1}{2} + \cos(\theta_\xi - \theta_x) + \cos 2(\theta_\xi - \theta_x) + \dots = 0,
\end{aligned}$$

and

$$\begin{aligned}
\underbrace{\arccos \xi}_{\theta_\xi} \neq -\underbrace{\arccos x}_{\theta_x} + 2\pi m &\Rightarrow \theta_\xi + \theta_x \neq 2\pi m \Rightarrow \\
&\Rightarrow \frac{1}{2} + \cos(\theta_\xi + \theta_x) + \cos 2(\theta_\xi + \theta_x) + \dots = 0.
\end{aligned}$$

Therefore, for $\xi \neq x$,

$$\frac{1}{2}T_0(\xi)T_0(x) + T_1(\xi)T_1(x) + T_2(\xi)T_2(x) + \dots = 0.$$

For $\xi = x$, there are two cases

case 1 $\underbrace{\arccos \xi}_{\theta_\xi} = \underbrace{\arccos x}_{\theta_x} + 2\pi m \Rightarrow \theta_\xi - \theta_x = 2\pi m \Rightarrow$
 $\Rightarrow \frac{1}{2} + \cos(\theta_\xi - \theta_x) + \cos 2(\theta_\xi - \theta_x) + \dots = \frac{1}{2} + 1 + 1 + \dots,$

If we also have $\theta_\xi = 2\pi k$, or $\theta_x = 2\pi k$, then $\theta_\xi + \theta_x = 2\pi l$, and

$$\frac{1}{2} + \cos(\theta_\xi + \theta_x) + \cos 2(\theta_\xi + \theta_x) + \dots = \frac{1}{2} + 1 + 1 + \dots$$

Then, the two cosine series sum up to

$$\frac{1}{2} T_0^2(x) + T_1^2(x) + T_2^2(x) + \dots = \frac{1}{2} + 1 + 1 + \dots$$

Else, $\frac{1}{2} + \cos(\theta_\xi + \theta_x) + \cos 2(\theta_\xi + \theta_x) + \dots$ vanishes, and

$$\frac{1}{2} T_0^2(x) + T_1^2(x) + T_2^2(x) + \dots = \frac{1}{2}(\frac{1}{2} + 1 + 1 + \dots).$$

case 2 $\underbrace{\arccos \xi}_{\theta_\xi} = -\underbrace{\arccos x}_{\theta_x} + 2\pi m \Rightarrow \theta_\xi + \theta_x = 2\pi m \Rightarrow$
 $\Rightarrow \frac{1}{2} + \cos(\theta_\xi + \theta_x) + \cos 2(\theta_\xi + \theta_x) + \dots = \frac{1}{2} + 1 + 1 + \dots.$

If we also have $\theta_\xi = 2\pi k$, or $\theta_x = 2\pi k$, then $\theta_\xi - \theta_x = 2\pi l$, and

$$\frac{1}{2} + \cos(\theta_\xi - \theta_x) + \cos 2(\theta_\xi - \theta_x) + \dots = \frac{1}{2} + 1 + 1 + \dots$$

Then, the two cosine series sum up to

$$\frac{1}{2} T_0^2(x) + T_1^2(x) + T_2^2(x) + \dots = \frac{1}{2} + 1 + 1 + \dots$$

Else, $\frac{1}{2} + \cos(\theta_\xi - \theta_x) + \cos 2(\theta_\xi - \theta_x) + \dots$ vanishes, and

$$\frac{1}{2} T_0^2(x) + T_1^2(x) + T_2^2(x) + \dots = \frac{1}{2}(\frac{1}{2} + 1 + 1 + \dots).$$

Therefore, for $\xi = x$,

$$\frac{1}{2}T_0^2(x) + T_1^2(x) + T_2^2(x) + \dots \geq \frac{1}{2}(1 + 1 + 1 + \dots)$$

Hence, the Chebyshev Kernel diverges to ∞ at any $\xi = x$.

Therefore, while the partial sums of the Chebyshev Series exist, their limit does not. That is, due to the singularity at $\xi = x$, the Chebyshev Series does not converge in the Calculus of Limits.

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because at any $\xi \neq x$, the Chebyshev Kernel vanishes, and the integral will be identically zero, for any function $f(x)$.


Plots of the Chebyshev Sequence confirm that

In the Calculus of Limits,

the Chebyshev Kernel is either singular or zero


1.2 Plots of $\frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\}$

In Maple, plot($\sum_{i=0}^{111} \frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \text{ChebyshevT}(i, 0.1) * \text{ChebyshevT}(i, x)$, $x = -.99 \dots .99$)




The pulse narrows with more terms:

In Maple, $\text{plot}\left(\sum_{i=0}^{365} \frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \text{ChebyshevT}(i, 0.2) * \text{ChebyshevT}(i, x), x = -.9 \dots .9\right)$




For $\theta = \arccos(x)$, the pulses are periodic

In Maple, $\text{plot}\left(\sum_{i=0}^{222} \frac{2}{\pi} \frac{1}{\sqrt{1-\cos^2(\theta)}} \text{ChebyshevT}(i, \cos(2)) * \text{ChebyshevT}(i, \cos(\theta)), \theta = -7 .. 7\right)$




In Maple, $\text{plot}\left(\sum_{i=0}^{222} \frac{2}{\pi} \frac{1}{\sqrt{1-\cos^2(\theta)}} \text{ChebyshevT}(i, \cos(0)) * \text{ChebyshevT}(i, \cos(\theta)), \theta = -13 .. .13\right)$



The computation is faster with the $T_n(x)$ written explicitly

In Maple $\text{plot}\left(\sum_{i=0}^{222} \frac{2}{\pi} \cdot \frac{1}{|\sin(\theta)|} \cdot \cos(i \cdot \theta), \theta = -37 \dots 37\right)$



The plots confirm that the Chebyshev Series Theorem cannot be proved in the Calculus of Limits.

1.3 Infinitesimal Calculus Solution

By resolving the problem of the infinitesimals [Dan2], we obtained the Infinite Hyper-reals that are strictly smaller than ∞ , and constitute the value of the Delta Function at the singularity.

The controversy surrounding the Leibnitz Infinitesimals derailed the development of the Infinitesimal Calculus, and the Delta Function could not be defined and investigated properly.

In Infinitesimal Calculus, [Dan3], we can differentiate over jump

discontinuities, and integrate over singularities.

The Delta Function, the idealization of an impulse in Radar circuits, is a Discontinuous Hyper-Real function which definition requires Infinite Hyper-reals, and which analysis requires Infinitesimal Calculus.

In [Dan5], we show that in infinitesimal Calculus, the hyper-real

$$\delta(x) = \frac{1}{2\pi} \int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega$$

is zero for any $x \neq 0$,

it spikes at $x = 0$, so that its Infinitesimal Calculus

integral is
$$\int_{x=-\infty}^{x=\infty} \delta(x) dx = 1,$$

and
$$\delta(0) = \frac{1}{dx} < \infty.$$

Here, we show that in Infinitesimal calculus, the Chebyshev Kernel is a periodic hyper-real Delta Function: A periodic train of Delta Functions.

And the Chebyshev Series $\mathcal{C}_{chebyshev} \mathcal{S}\{f(x)\}$ associated with a Hyper-real periodic function $f(x)$, equals $f(x)$.

2.

Hyper-real Line

Each real number α can be represented by a Cauchy sequence of rational numbers, (r_1, r_2, r_3, \dots) so that $r_n \rightarrow \alpha$.

The constant sequence $(\alpha, \alpha, \alpha, \dots)$ is a constant hyper-real.

In [Dan2] we established that,

1. Any totally ordered set of positive, monotonically decreasing to zero sequences (l_1, l_2, l_3, \dots) constitutes a family of infinitesimal hyper-reals.
2. The infinitesimals are smaller than any real number, yet strictly greater than zero.
3. Their reciprocals $\left(\frac{1}{l_1}, \frac{1}{l_2}, \frac{1}{l_3}, \dots\right)$ are the infinite hyper-reals.
4. The infinite hyper-reals are greater than any real number, yet strictly smaller than infinity.
5. The infinite hyper-reals with negative signs are smaller than any real number, yet strictly greater than $-\infty$.
6. The sum of a real number with an infinitesimal is a non-constant hyper-real.
7. The Hyper-reals are the totality of constant hyper-reals, a family of infinitesimals, a family of infinitesimals with

- negative sign, a family of infinite hyper-reals, a family of infinite hyper-reals with negative sign, and non-constant hyper-reals.
8. The hyper-reals are totally ordered, and aligned along a line: the Hyper-real Line.
 9. That line includes the real numbers separated by the non-constant hyper-reals. Each real number is the center of an interval of hyper-reals, that includes no other real number.
 10. In particular, zero is separated from any positive real by the infinitesimals, and from any negative real by the infinitesimals with negative signs, $-dx$.
 11. Zero is not an infinitesimal, because zero is not strictly greater than zero.
 12. We do not add infinity to the hyper-real line.
 13. The infinitesimals, the infinitesimals with negative signs, the infinite hyper-reals, and the infinite hyper-reals with negative signs are semi-groups with respect to addition. Neither set includes zero.
 14. The hyper-real line is embedded in \mathbb{R}^∞ , and is not homeomorphic to the real line. There is no bi-continuous one-one mapping from the hyper-real onto the real line.

15. In particular, there are no points on the real line that can be assigned uniquely to the infinitesimal hyper-reals, or to the infinite hyper-reals, or to the non-constant hyper-reals.
16. No neighbourhood of a hyper-real is homeomorphic to an \mathbb{R}^n ball. Therefore, the hyper-real line is not a manifold.
17. The hyper-real line is totally ordered like a line, but it is not spanned by one element, and it is not one-dimensional.

3.

Integral of a Hyper-real Function

In [Dan3], we defined the integral of a Hyper-real Function.

Let $f(x)$ be a hyper-real function on the interval $[a, b]$.

The interval may not be bounded.

$f(x)$ may take infinite hyper-real values, and need not be bounded.

At each

$$a \leq x \leq b,$$

there is a rectangle with base $[x - \frac{dx}{2}, x + \frac{dx}{2}]$, height $f(x)$, and area

$$f(x)dx.$$

We form the **Integration Sum** of all the areas for the x 's that start at $x = a$, and end at $x = b$,

$$\sum_{x \in [a, b]} f(x)dx.$$

If for any infinitesimal dx , the Integration Sum has the same hyper-real value, then $f(x)$ is integrable over the interval $[a, b]$.

Then, we call the Integration Sum the integral of $f(x)$ from $x = a$, to $x = b$, and denote it by

$$\int_{x=a}^{x=b} f(x)dx .$$

If the hyper-real is infinite, then it is the integral over $[a, b]$,

If the hyper-real is finite,

$$\int_{x=a}^{x=b} f(x)dx = \text{real part of the hyper-real. } \square$$

3.1 The countability of the Integration Sum

In [Dan1], we established the equality of all positive infinities:

We proved that the number of the Natural Numbers,

$Card\mathbb{N}$, equals the number of Real Numbers, $Card\mathbb{R} = 2^{Card\mathbb{N}}$, and

we have

$$Card\mathbb{N} = (Card\mathbb{N})^2 = \dots = 2^{Card\mathbb{N}} = 2^{2^{Card\mathbb{N}}} = \dots \equiv \infty .$$

In particular, we demonstrated that the real numbers may be well-ordered.

Consequently, there are countably many real numbers in the interval $[a, b]$, and the Integration Sum has countably many terms.

While we do not sequence the real numbers in the interval, the summation takes place over countably many $f(x)dx$.

The Lower Integral is the Integration Sum where $f(x)$ is replaced

by its lowest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.2} \quad \sum_{x \in [a, b]} \left(\inf_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

The Upper Integral is the Integration Sum where $f(x)$ is replaced

by its largest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.3} \quad \sum_{x \in [a, b]} \left(\sup_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

If the integral is a finite hyper-real, we have

3.4 *A hyper-real function has a finite integral if and only if its upper integral and its lower integral are finite, and differ by an infinitesimal.*

4.

Delta Function

In [Dan5], we have defined the Delta Function, and established its properties

1. The Delta Function is a hyper-real function defined from the

hyper-real line into the set of two hyper-reals $\left\{0, \frac{1}{dx}\right\}$. The

hyper-real 0 is the sequence $\langle 0, 0, 0, \dots \rangle$. The infinite hyper-

real $\frac{1}{dx}$ depends on our choice of dx .

2. We will usually choose the family of infinitesimals that is

spanned by the sequences $\left\langle \frac{1}{n} \right\rangle, \left\langle \frac{1}{n^2} \right\rangle, \left\langle \frac{1}{n^3} \right\rangle, \dots$. It is a

semigroup with respect to vector addition, and includes all

the scalar multiples of the generating sequences that are

non-zero. That is, the family includes infinitesimals with

negative sign. Therefore, $\frac{1}{dx}$ will mean the sequence $\langle n \rangle$.

Alternatively, we may choose the family spanned by the

sequences $\left\langle \frac{1}{2^n} \right\rangle, \left\langle \frac{1}{3^n} \right\rangle, \left\langle \frac{1}{4^n} \right\rangle, \dots$. Then, $\frac{1}{dx}$ will mean the

sequence $\langle 2^n \rangle$. Once we determined the basic infinitesimal dx , we will use it in the Infinite Riemann Sum that defines an Integral in Infinitesimal Calculus.

3. The Delta Function is strictly smaller than ∞

4. We define, $\delta(x) \equiv \frac{1}{dx} \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x)$,

$$\text{where } \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x) = \begin{cases} 1, & x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right] \\ 0, & \text{otherwise} \end{cases}.$$

5. Hence,

$$\diamond \text{ for } x < 0, \delta(x) = 0$$

$$\diamond \text{ at } x = -\frac{dx}{2}, \delta(x) \text{ jumps from } 0 \text{ to } \frac{1}{dx},$$

$$\diamond \text{ for } x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right], \delta(x) = \frac{1}{dx}.$$

$$\diamond \text{ at } x = 0, \delta(0) = \frac{1}{dx}$$

$$\diamond \text{ at } x = \frac{dx}{2}, \delta(x) \text{ drops from } \frac{1}{dx} \text{ to } 0.$$

$$\diamond \text{ for } x > 0, \delta(x) = 0.$$

$$\diamond x\delta(x) = 0$$

6. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \left\langle \mathcal{X}_{[-\frac{1}{2}, \frac{1}{2}]}(x), 2\mathcal{X}_{[-\frac{1}{4}, \frac{1}{4}]}(x), 3\mathcal{X}_{[-\frac{1}{6}, \frac{1}{6}]}(x), \dots \right\rangle$

7. If $dx = \langle \frac{2}{n} \rangle$, $\delta(x) = \left\langle \frac{1}{2 \cosh^2 x}, \frac{2}{2 \cosh^2 2x}, \frac{3}{2 \cosh^2 3x}, \dots \right\rangle$

8. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \langle e^{-x} \chi_{[0, \infty)}, 2e^{-2x} \chi_{[0, \infty)}, 3e^{-3x} \chi_{[0, \infty)}, \dots \rangle$

9.
$$\int_{x=-\infty}^{x=\infty} \delta(x) dx = 1.$$

10.
$$\delta(\xi - x) = \frac{1}{2\pi} \int_{k=-\infty}^{k=\infty} e^{-ik(\xi-x)} dk$$

5.

Periodic Delta $\delta_{Periodic}(\theta_\xi - \theta_x)$

5.1 Periodic Delta Function Definition

$$\delta_{Periodic}(\theta_\xi - \theta_x) = \dots + \delta(\theta_\xi - \theta_x + 2\pi) + \delta(\theta_\xi - \theta_x) + \delta(\theta_\xi - \theta_x - 2\pi) + \dots$$

is a periodic hyper-real Delta function, with period $T = 2\pi$.

5.2 Fourier Transform of $\delta_{Periodic}(\theta)$

$$\mathcal{F}\{\delta_{Periodic}(\theta)\} = \dots + e^{-i4\pi^2\nu} + 1 + e^{i4\pi^2\nu} + \dots$$

Proof: $\mathcal{F}\{\delta_{Periodic}(\theta)\} = \dots + \mathcal{F}\{\delta(\theta + 2\pi)\} + \mathcal{F}\{\delta(\theta)\} + \mathcal{F}\{\delta(\theta - 2\pi)\} + \dots$

$$= \dots + \underbrace{\int_{\theta=-\infty}^{\theta=\infty} \delta(\theta + 2\pi)e^{-i2\pi\nu\theta} d\theta}_{e^{i2\pi 2\pi\nu}} + \underbrace{\int_{\theta=-\infty}^{\theta=\infty} \delta(\theta)e^{-i2\pi\nu\theta} d\theta}_1 + \underbrace{\int_{\theta=-\infty}^{\theta=\infty} \delta(\theta - 2\pi)e^{-i2\pi\nu\theta} d\theta}_{e^{-i2\pi 2\pi\nu}} + \dots$$

5.3 Fourier Integral Theorem for $\delta_{Periodic}(\theta)$

$$\mathcal{F}^{-1}\mathcal{F}\{\delta_{Periodic}(\theta)\} = \delta_{Periodic}(\theta)$$

Proof: $\mathcal{F}^{-1}\mathcal{F}\{\delta_{Periodic}(\theta)\} = \dots + \mathcal{F}^{-1}\{e^{2\pi i 2\pi\nu}\} + \mathcal{F}^{-1}\{1\} + \mathcal{F}^{-1}\{e^{-2\pi i 2\pi\nu}\} + \dots$

$$= \dots + \underbrace{\int_{\nu=-\infty}^{\nu=\infty} e^{i2\pi\nu 2\pi} e^{i2\pi\nu\theta} d\nu}_{\delta(\theta+2\pi)} + \underbrace{\int_{\nu=-\infty}^{\nu=\infty} e^{i2\pi\nu\theta} d\nu}_{\delta(\theta)} + \underbrace{\int_{\nu=-\infty}^{\nu=\infty} e^{-i2\pi\nu 2\pi} e^{i2\pi\nu\theta} d\nu}_{\delta(\theta-2\pi)} + \dots \square$$

6.

Convergent Series

In [Dan8], we defined convergence of infinite series in Infinitesimal Calculus

6.1 Sequence Convergence to a finite hyper-real a

$$a_n \rightarrow a \text{ iff } \langle a_n \rangle - \langle a \rangle = \text{infinitesimal.}$$

6.2 Sequence Convergence to an infinite hyper-real A

$$a_n \rightarrow A \text{ iff } \langle a_n \rangle \text{ represents the infinite hyper-real } A.$$

6.3 Series Convergence to a finite hyper-real s

$$a_1 + a_2 + \dots \rightarrow s \text{ iff } \langle a_1 + \dots + a_n \rangle - \langle s \rangle = \text{infinitesimal.}$$

6.4 Series Convergence to an Infinite Hyper-real S

$$a_1 + a_2 + \dots \rightarrow S \text{ iff}$$

$$\langle a_1 + \dots + a_n \rangle \text{ represents the infinite hyper-real } S.$$

7.

Chebyshev Sequence and $\delta_{Periodic}(\theta_\xi - \theta_x)$

7.1 Chebyshev Sequence Definition

The Chebyshev Series partial sums

$$C_{chebyshev} \mathcal{S}_n \{f(x)\} = \int_{\xi=-1}^{\xi=1} f(\xi) \underbrace{\frac{2}{\pi \sqrt{1-x^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\}}_{\text{Chebyshev Sequence}} d\xi,$$

give rise to the Chebyshev Sequence

$$T_n(\xi, x) = \frac{2}{\pi \sqrt{1-x^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\}.$$

7.2 Chebyshev Sequence is a Periodic Delta Sequence

For each $n = 0, 1, 2, 3, \dots$,

$$\begin{aligned} T_n(\xi, x) &= \frac{2}{\pi \sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\}, \\ &= \frac{2}{\pi \sqrt{1-\xi^2}} \left\{ \frac{1}{2} + \cos \theta_\xi \cos \theta_x + \dots + \cos n\theta_\xi \cos n\theta_x \right\} \end{aligned}$$

1. *has the sifting property on each interval,*

$$\dots \int_{\xi=-3}^{\xi=-1} T_n(\xi, x) d\xi = 1; \quad \int_{\xi=-1}^{\xi=1} T_n(\xi, x) d\xi = 1; \quad \int_{\xi=1}^{\xi=3} T_n(\xi, x) d\xi = 1 \dots$$

2. *is a continuous function*

3. *peaks on each of these intervals to* $\lim_{\xi \rightarrow x} T_n(\xi, x) \geq \frac{1}{2}(n + 1)$

Proof of (1)

$$\begin{aligned} & \int_{\xi=-1}^{\xi=1} \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\} d\xi = \\ &= \frac{1}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} d\xi + T_1(x) \frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} T_1(\xi) d\xi + \dots \\ & \qquad \qquad \qquad \dots + T_n(x) \frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} T_n(\xi) d\xi \end{aligned}$$

The first integral transforms with $\xi = \cos \theta$, $d\xi = -\sin \theta d\theta$, to

$$\frac{1}{\pi} \int_{\theta=\pi}^{\theta=0} \frac{1}{\sqrt{1-\cos^2 \theta}} (-\sin \theta) d\theta = \frac{1}{\pi} \int_{\theta=0}^{\theta=\pi} d\theta = 1.$$

The terms with $k = 1, 2, \dots, n$, vanish because of the orthogonality of the Chebyshev Polynomials:

By [Spiegel, p.156, #30.18], for $k = 1, 2, \dots, n$,

$$\int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} T_k(\xi) d\xi = \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} T_0(\xi) T_k(\xi) d\xi = \delta_{0k} = 0.$$

Thus, for $n = 0, 1, 2, \dots$, $\int_{\xi=-1}^{\xi=1} \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\} d\xi = 1. \square$

Proof of (3) in 1.1. \square

8.

Chebyshev Kernel and $\delta_{Periodic}(\theta_\xi - \theta_x)$

8.1 Chebyshev Kernel in the Calculus of Limits

The Chebyshev Series partial sums

$$\mathcal{C}_{hebyshev} \mathcal{S}_n \{f(x)\} = \int_{\xi=-1}^{\xi=1} f(\xi) \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \underbrace{\left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) \right\}}_{\text{Chebyshev Sequence}} d\xi.$$

give rise to the Chebyshev Sequence.

The limit of the Chebyshev Sequence is an infinite series, the Chebyshev Kernel

$$\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) + \dots \right\}$$

8.2 *In the Calculus of Limits, the Chebyshev Kernel does not have the sifting property*

Proof: for $\xi \rightarrow x$, $\frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \left\{ \frac{1}{2} + T_1^2(x) + \dots + T_n^2(x) \right\} \geq \frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \frac{1}{2} (n+1)$

$$\xrightarrow[n \rightarrow \infty]{} \infty$$

That is, $\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) + \dots \right\}$ is singular,

and cannot be integrated over. \square

8.3 Hyper-real Chebyshev Kernel in Infinitesimal Calculus

Let $x = \cos \theta_x$, $\xi = \cos \theta_\xi$, $\langle n \rangle$ an infinite Hyper-real. Then

$$\begin{aligned}
 \mathcal{C}_{hebyshev}(\theta_\xi - \theta_x) &= \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) + \dots \right\} \\
 &= \begin{cases} \langle n \rangle, & \theta_\xi - \theta_x = 2\pi m \\ 0, & \theta_\xi - \theta_x \neq 2\pi m \end{cases} \\
 &= \dots + \delta(\theta_\xi - \theta_x + 2\pi) + \delta(\theta_\xi - \theta_x) + \delta(\theta_\xi - \theta_x - 2\pi) + \dots \\
 &= \delta_{Periodic}(\theta_\xi - \theta_x).
 \end{aligned}$$

Proof: $\mathcal{C}_{hebyshev}(\theta_\xi - \theta_x) = \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots + T_n(\xi)T_n(x) + \dots \right\}$

$$\begin{aligned}
 &= \begin{cases} \langle n \rangle, & \theta_\xi - \theta_x = 2\pi m \\ 0, & \theta_\xi - \theta_x \neq 2\pi m \end{cases} \\
 &= \dots + \begin{cases} 0, \theta_\xi - \theta_x \neq -2\pi \\ \frac{1}{d\theta}, \theta_\xi - \theta_x = -2\pi \end{cases} + \begin{cases} 0, \theta_\xi - \theta_x \neq 0 \\ \frac{1}{d\theta}, \theta_\xi - \theta_x = 0 \end{cases} + \begin{cases} 0, \theta_\xi - \theta_x \neq 2\pi \\ \frac{1}{d\theta}, \theta_\xi - \theta_x = 2\pi \end{cases} + \dots \\
 &= \dots + \delta(\theta_\xi - \theta_x + 2\pi) + \delta(\theta_\xi - \theta_x) + \delta(\theta_\xi - \theta_x - 2\pi) + \dots \\
 &= \delta_{Periodic}(\theta_\xi - \theta_x). \square
 \end{aligned}$$

9.

Chebyshev Series and $\delta_{Periodic}(\theta_\xi - \theta_x)$

9.1 Chebyshev Series of a Hyper-real Function

Let $f(x)$ be a hyper-real function integrable on $[-1,1]$.

For $x = \cos \theta_x$, $f(\cos \theta_x)$ is defined for each real θ_x .

Then, for each $n = 0,1,2,3,\dots$, the integrals

$$a_n = \frac{2}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} f(x) T_n(x) dx$$

exist, with finite, or infinite hyper-real values. The a_n are the Chebyshev Series Coefficients of $f(x)$.

The Chebyshev Series associated with $f(x)$ is

$$\begin{aligned} \mathcal{C}_{chebyshev} \mathcal{S} \{ f(x) \} &= a_0 \frac{1}{2} + a_1 T_1(x) + a_2 T_2(x) + a_3 T_3(x) + \dots \\ &= a_0 \frac{1}{2} + a_1 \cos \theta_x + a_2 \cos 2\theta_x + a_3 \cos n\theta_x + \dots \end{aligned}$$

For each x , it may assume finite or infinite hyper-real values.

$$\mathbf{9.2} \quad \mathcal{C}_{chebyshev} \mathcal{S} \{ \delta_{Periodic}(\theta_\xi - \theta_x) \} = \delta_{Periodic}(\theta_\xi - \theta_x)$$

Proof:

$$\mathcal{C}_{chebyshev} \mathcal{S} \{ \delta_{Periodic}(\theta_\xi - \theta_x) \} = a_0 + a_1 T_1(x) + a_2 T_2(x) + a_3 T_3(x) \dots$$

where

$$a_n = \frac{2}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} \delta_{Periodic}(\theta_\xi - \theta_x) T_n(x) dx,$$

Substituting from 8.3,

$$\delta_{Periodic}(\theta_\xi - \theta_x) = \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) + \dots T_m(\xi) T_m(x) + \dots \right\},$$

$$\begin{aligned} a_n &= \frac{2}{\pi} \int_{x=-1}^{x=1} \frac{1}{\sqrt{1-x^2}} \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) + \dots T_m(\xi) T_m(x) + \dots \right\} T_n(x) dx \\ &= \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} T_0(\xi) \underbrace{\int_{x=-1}^{x=1} \frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \frac{1}{2} T_0(x) T_n(x) dx}_{0} + \dots \\ &\quad \dots + \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} T_m(\xi) \underbrace{\int_{x=-1}^{x=1} \frac{2}{\pi} \frac{1}{\sqrt{1-x^2}} \frac{1}{2} T_m(x) T_n(x) dx}_{\delta_{mn}} + \dots \\ &= \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} T_n(\xi). \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{L}_{Legendre} \mathcal{S} \left\{ \delta_{Periodic}(\theta_\xi - \theta_x) \right\} &= \\ &= \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) + \dots T_m(\xi) T_m(x) + \dots \right\} \\ &= \delta_{Periodic}(\theta_\xi - \theta_x). \square \end{aligned}$$

10.

Chebyshev Series Theorem

The Chebyshev Series Theorem for a hyper-real function, $f(x)$, is the Fundamental Theorem of Chebyshev Series.

It supplies the conditions under which the Chebyshev Series associated with $f(x)$ equals $f(x)$.

It is believed to hold in the Calculus of Limits. In fact,

The Theorem cannot be proved in the Calculus of Limits under any conditions,

because the summation of the Chebyshev Series requires integration of the singular Chebyshev Kernel.

10.1 Chebyshev Series Theorem cannot be proved in the Calculus of Limits

Proof: Let $f(x)$ be integrable on $[-1,1]$, so that $f(1) = f(-1)$

In the Calculus of Limits, the Chebyshev Series is the limit of

$$\begin{aligned} \mathcal{C}_{chebyshev} \mathcal{S}_n \{f(x)\} &= \frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + \dots + a_n T_n(x) \\ &= \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_0(\xi) d\xi \right) T_0(x) + \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_1(\xi) d\xi \right) T_1(x) + \dots \end{aligned}$$

$$\dots + \left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_0(\xi) d\xi \right) T_n(x)$$

$$= \int_{\xi=-1}^{\xi=1} f(\xi) \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) \dots + T_n(\xi) T_n(x) \right\} d\xi.$$

As $n \rightarrow \infty$, the Chebyshev Sequence

$$\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) \dots + T_n(\xi) T_n(x) \right\}$$

becomes the Chebyshev Kernel,

$$\frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} T_0(\xi) T_0(x) + T_1(\xi) T_1(x) \dots + T_n(\xi) T_n(x) + \dots \right\},$$

By 8.2, The Chebyshev Kernel diverges to infinity at any $\xi = x$.

In particular,

$$x = \xi \Rightarrow \theta_\xi = \theta_x + 2\pi m,$$

and the Chebyshev Kernel diverges to infinity.

Therefore, while the partial sums of the Chebyshev Series exist, their limit does not. The sifting through the values of $f(\xi)$ by the Chebyshev Kernel, and the picking of $f(\xi)$ at $\xi = x$, is unrecognized in the Calculus of Limits

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because for any $\xi \neq x$, the Chebyshev Kernel vanishes, and the integral is identically zero, for any function $f(x)$.

Thus, the Chebyshev Series Theorem cannot be proved in the Calculus of Limits. \square

10.2 Calculus of Limits Conditions are irrelevant to Chebyshev Series Theorem

Proof: It is clear from 10.1 that Calculus of Limits conditions requiring smoothness of $f(x)$ will not resolve the singularity of the Chebyshev kernel, and are not sufficient for the Chebyshev Series Theorem. \square

In Infinitesimal Calculus, by 8.3, the Chebyshev Kernel is the Periodic Delta Function, and by 9.2, it equals its Chebyshev Series.

Then, the Chebyshev Series Theorem holds for any periodic Hyper-Real Function:

10.3 Chebyshev Series Theorem for periodic Hyper-real

$f(x)$

If $f(x)$ is hyper-real function integrable on $[-1,1]$, and $f(-1) = f(1)$

Then,

$$f(x) = C_{chebyshev} \mathcal{S} \{ f(x) \}$$

Proof:

$$f(x) = \int_{\xi=-1}^{\xi=1} f(\xi) \underbrace{\left\{ \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots \right\}}_{\delta_{\text{Periodic}}(\xi-x), \text{ where the period of Delta is 2}} d\xi$$

Substituting from 8.3,

$$\delta_{\text{Periodic}}(\xi - x) = \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots T_m(\xi)T_m(x) + \dots \right\},$$

$$f(x) = \int_{\xi=-1}^{\xi=1} f(\xi) \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots T_m(\xi)T_m(x) + \dots \right\} d\xi$$

This Hyper-real Integral is the summation,

$$\sum_{\xi=-1}^{\xi=1} f(\xi) \frac{2}{\pi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \frac{1}{2} + T_1(\xi)T_1(x) + \dots T_m(\xi)T_m(x) + \dots \right\} d\xi$$

which amounts to the hyper-real function $f(x)$, and is well-defined.

Hence, the summation of each term in the integrand exists, and we may write the integral as the sum

$$= \frac{1}{2} \underbrace{\left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_0(\xi) d\xi \right)}_{\frac{1}{2}a_0} \underbrace{T_0(x)}_1 + \underbrace{\left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_1(\xi) d\xi \right)}_{a_1} T_1(x) + \dots$$

$$\dots + \underbrace{\left(\frac{2}{\pi} \int_{\xi=-1}^{\xi=1} \frac{1}{\sqrt{1-\xi^2}} f(\xi) T_n(\xi) d\xi \right)}_{a_n} T_n(x) + \dots$$

$$= \frac{1}{2} a_0 T_0(x) + a_1 T_1(x) + a_2 T_2(x) + \dots$$

$$= \mathcal{C}_{\text{hebyshhev}} \mathcal{S} \{ f(x) \}. \square$$

In particular, the periodic Delta Function is not smooth.

❖ *The Hyper-real $\delta(x)$, is not defined in the Calculus of Limits,
and is not integrable in $[-1,1]$.*

But by 9.2, $\delta_{Periodic}(x)$ satisfies the Chebyshev Series Theorem.

References

[Abramowitz] Abramowitz, M., and Stegun, I., “*Handbook of Mathematical Functions with Formulas Graphs and Mathematical Tables*”, U.S. Department of Commerce, National Bureau of Standards, 1964.

[Dan1] Dannon, H. Vic, “*Well-Ordering of the Reals, Equality of all Infinities, and the Continuum Hypothesis*” in Gauge Institute Journal Vol. 6 No. 2, May 2010;

[Dan2] Dannon, H. Vic, “*Infinitesimals*” in Gauge Institute Journal Vol.6 No. 4, November 2010;

[Dan3] Dannon, H. Vic, “*Infinitesimal Calculus*” in Gauge Institute Journal Vol. 7 No. 4, November 2011;

[Dan4] Dannon, H. Vic, “*Riemann’s Zeta Function: the Riemann Hypothesis Origin, the Factorization Error, and the Count of the Primes*”, in Gauge Institute Journal of Math and Physics, Vol. 5, No. 4, November 2009.

[Dan5] Dannon, H. Vic, “*The Delta Function*” in Gauge Institute Journal Vol. 8, No. 1, February, 2012;

[Dan6] Dannon, H. Vic, “*Riemannian Trigonometric Series*”, Gauge Institute Journal, Volume 7, No. 3, August 2011.

[Dan7] Dannon, H. Vic, “*Delta Function the Fourier Transform, and the Fourier Integral Theorem*” in Gauge Institute Journal Vol. 8, No. 2, May, 2012;

[Dan8] Dannon, H. Vic, “*Infinite Series with Infinite Hyper-real Sum* ” in Gauge Institute Journal Vol. 8, No. 3, August, 2012;

[Ferrers] Ferrers, N., M., “*An Elementary treatment on Spherical Harmonics*”, Macmillan, 1877.

[Gradshteyn] Gradshteyn, I., S., and Ryzhik, I., M., “*Tables of Integrals Series and Products*”, 7th Edition, edited by Allan Jeffery, and Daniel Zwillinger, Academic Press, 2007

[Hardy] Hardy, G. H., *Divergent Series*, Chelsea 1991.

[Krylov] Krylov, V. I., “*Approximate Calculation of Integrals*” Macmillan, 1962.

[Magnus] Magnus, W., Oberhettinger, F., Sony, R., P., “*Formulas and Theorems for the Special Functions of Mathematical Physics*” Third Edition, Springer-Verlag, 1966.

[Spanier] Spanier, Jerome, and Oldham, Keith, “*An Atlas of Functions*”, Hemisphere, 1987.

[Spiegel] Spiegel, Murray, “*Mathematical Handbook of formulas and tables*” Schaum’s Outline Series, McGraw Hill, 1968.

[Weisstein], Weisstein, Eric, W., “*CRC Encyclopedia of Mathematics*”, Third Edition, CRC Press, 2009.