

Fourier Transforms, and Fourier Integrals of a Complex Delta Function, and of an Analytic Function

H. Vic Dannon

vic0@comcast.net

January, 2011

Revised December, 2014

Abstract In the Calculus of Limits, it is erroneously believed [Dan4] that the existence of the Fourier Transform requires Fourier boundedness conditions on the function.

In fact, these conditions do not guarantee the existence of the Fourier Transform, and are irrelevant to the Transform's existence [Dan4].

When these unnecessary conditions are imposed on a function of complex variable, they are believed to guarantee the existence of the Fourier Transform of $f(z)$ along a path parallel to the x axis [Titchmarsh, p.44].

Then, the Fourier Transform is an analytic function in a strip parallel to the x axis. At the borders of the strip, the Analytic

Function which is inseparable from its disk of convergence, will have one sided derivative.

No one, including Titchmarsh, cared to explore this unnatural geometry, and the Fourier Transform that is analytic in a strip parallel to the x axis, was never developed further.

We define here the Fourier Transform of the Complex Delta Function, and the Fourier Transform of a hyper-complex Analytic function $f(z)$, along a closed path γ_ζ in the complex plane ζ .

Then,

1. The Hyper-complex Delta is given by its Fourier Integral

$$\delta(\zeta - z) = \frac{1}{2\pi} \oint_{|\omega|=\eta} e^{i\omega(\zeta-z)} d\omega = \frac{1}{2\pi} \oint_{|\omega|=1} e^{i\omega(\zeta-z)} d\omega,$$

where η is an infinitesimal.

2. For an Analytic Function $f(z)$, the Cauchy Integral Formula

$$f(z) = \frac{1}{2\pi i} \oint_{\gamma_\zeta} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

is the Fourier Integral Theorem

$$f(z) = \frac{1}{2\pi} \oint_{\gamma_\omega} \left(\oint_{\gamma_\zeta} f(\zeta) e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega.$$

3. The convergence of the Fourier Integral for an Analytic

Function implies the existence of the Fourier Transform of $f(z)$, and the existence of its inverse transform:

If $f(z)$ is a hyper-complex analytic function on a hyper-complex domain that includes the circle $|\zeta - z| = 1$,

Then,

1) the hyper-complex integral

$$\oint_{|\zeta-z|=1} f(\zeta)e^{-i\omega\zeta}d\zeta$$

converges to $\hat{f}(\omega)$

2) the hyper-complex integral

$$\frac{1}{2\pi} \oint_{|\omega|=1} \hat{f}(\omega)e^{iz\omega}d\omega$$

converges to $f(z)$

Keywords: Infinitesimal, Infinite-Hyper-Real, Hyper-Real, Cardinal, Infinity. Non-Archimedean, Non-Standard Analysis, Calculus, Limit, Continuity, Derivative, Integral, Complex Variable, Complex Analysis, Analytic Functions, Holomorphic, Cauchy Integral Theorem, Cauchy Integral Formula, Contour Integral.

2000 Mathematics Subject Classification 26E35; 26E30;
26E15; 26E20; 26A06; 26A12; 03E10; 03E55; 03E17; 03H15;
46S20; 97I40; 97I30.

Contents

Introduction

1. Hyper-real Line
2. Hyper-Complex Plane
3. Hyper-Complex Function
4. Hyper-Complex Path Integral
5. Hyper-real Delta Function
6. Hyper-Complex Delta Function $\delta(z)$
7. Cauchy Integral Formula
8. Hyper-real Fourier Transform
9. Fourier Transform of Hyper-Complex $f(z)$
10. Fourier Transform of $\delta(z)$
11. Fourier Integral of $\delta(z)$
12. Fourier Integral of an Analytic $f(z)$
13. Fourier Transform of an Analytic $f(z)$

References

Introduction

0.1 Complex Fourier Transform in the Calculus of Limits

In the Calculus of Limits, it is erroneously believed [Dan4] that the existence of the Fourier Transform requires Fourier boundedness conditions on the function.

When these unnecessary conditions are imposed on a function of complex variable, they are believed to guarantee the existence of the Fourier Transform of $f(z)$ along a path parallel to the x axis [Titchmarsh, p.44].

Then, the Fourier Transform is an analytic function in a strip parallel to the x axis. At the borders of the strip, the Analytic Function which is inseparable from its disk of convergence, will have one sided derivative.

In [Titchmarsh, p. 44], Titchmarsh has

THEOREM 26.

Let $f(z)$ ***be an analytic function, regular for***

$$-a < y < b, \quad \text{where } a > 0, b > 0.$$

In any strip interior to $-a < y < b$, ***and for any*** $\varepsilon > 0$, ***let***

$$f(z) = \begin{cases} O(e^{-(\lambda-\varepsilon)x}), & (x \rightarrow \infty) \\ O(e^{(\mu-\varepsilon)x}), & (x \rightarrow -\infty) \end{cases}'$$

where $\lambda > 0$, $\mu > 0$.

Then,

$$F(\omega) = \frac{1}{\sqrt{2\pi}} \int_{\zeta=-\infty}^{\zeta=\infty} f(\zeta)e^{i\omega\zeta} d\zeta$$

satisfies conditions similar to those imposed on $f(z)$,

with a, b, λ, μ replaced by λ, μ, b, a ; and

$$f(z) = \frac{1}{\sqrt{2\pi}} \int_{\omega=-\infty}^{\omega=\infty} F(\omega)e^{-i\omega z} d\omega$$

In fact, these conditions

- do not guarantee the existence of the Fourier Transform,
- are irrelevant to the Transform's existence [Dan4],
- apply to an Analytic Function in an unnatural domain of a strip, rather than a disk, miscomprehending analyticity in the complex plane.


No one, including Titchmarsh, cared to explore this unnatural geometry, and the Fourier Transform that is analytic in a strip parallel to the x axis, was never developed further.

0.2 Limitation of the Integration Path to the Real Line

The Conditions of Theorem 26 above, limit the integration path to the real line, and replace the powerful analysis of analytic

functions in the complex plane with limited power of analysis of differentiable functions on the line.

Instead of integrating along a closed path such as



which may be homologous to a unit circle, the Conditions require the function to approximately vanish out of a compact interval, and limit the admissible functions that may have a Fourier Transform.

For an Analytic Function these Conditions are unnecessary: We show here that the Cauchy Integral Formula, along a closed path, yields the Fourier Integral along that closed path. Therefore, the Fourier Integral converges along the arc from ∞ to $-\infty$, and along the real line from $-\infty$ to ∞ . Consequently, the Fourier Transform along the real line from $-\infty$ to ∞ converges, without any restrictions of exponential decay on the function.

The limitation is particularly awkward considering that as an Analytic function, the Fourier Transform $F(\omega)$ is inseparable from

its disk of convergence, and its derivative may be obtained in any radial direction. In Titchmarsh's presentation, at the borders of its strip of convergence, $F(\omega)$ has one sided derivative.

0.3 Definition along a Closed Path

Here, we show that for an Analytic Function, the Cauchy Integral Formula is the Fourier Integral of that Analytic Function.

That is, for an Analytic function, the Cauchy Integral Formula, and the Fourier Integral, coincide.

The Fourier Integral defines the Fourier Transform of an Analytic function $f(z)$ along a closed path.

Thus, for an Analytic Function, the Fourier Transform exists with no need for the Calculus of Limits Conditions of Titchmarsh's Theorem 26.

We start by recalling the Hyper-real line, and the Hyper-Complex Plane.

1.

Hyper-real Line

Each real number α can be represented by a Cauchy sequence of rational numbers, (r_1, r_2, r_3, \dots) so that $r_n \rightarrow \alpha$.

The constant sequence $(\alpha, \alpha, \alpha, \dots)$ is a constant Hyper-real.

In [Dan2] we established that,

1. Any totally ordered set of positive, monotonically decreasing to zero sequences (l_1, l_2, l_3, \dots) constitutes a family of infinitesimal Hyper-reals.
2. The infinitesimals are smaller than any real number, yet strictly greater than zero.
3. Their reciprocals $\left(\frac{1}{l_1}, \frac{1}{l_2}, \frac{1}{l_3}, \dots\right)$ are the infinite Hyper-reals.
4. The infinite Hyper-reals are greater than any real number, yet strictly smaller than infinity.
5. The infinite Hyper-reals with negative signs are smaller than any real number, yet strictly greater than $-\infty$.
6. The sum of a real number with an infinitesimal is a non-constant Hyper-real.

7. The Hyper-reals are the totality of constant Hyper-reals, a family of infinitesimals, a family of infinitesimals with negative sign, a family of infinite Hyper-reals, a family of infinite Hyper-reals with negative sign, and non-constant Hyper-reals.
8. The Hyper-reals are totally ordered, and aligned along a line: the Hyper-real Line.
9. That line includes the real numbers separated by the non-constant Hyper-reals. Each real number is the center of an interval of Hyper-reals, that includes no other real number.
10. In particular, zero is separated from any positive real by the infinitesimals, and from any negative real by the infinitesimals with negative signs, $-dx$.
11. Zero is not an infinitesimal, because zero is not strictly greater than zero.
12. We do not add infinity to the Hyper-real line.
13. The infinitesimals, the infinitesimals with negative signs, the infinite Hyper-reals, and the infinite Hyper-reals with negative signs are semi-groups with respect to addition. Neither set includes zero.

14. The Hyper-real line is embedded in \mathbb{R}^∞ , and is not homeomorphic to the real line. There is no bi-continuous one-one mapping from the Hyper-real onto the real line.
15. In particular, there are no points on the real line that can be assigned uniquely to the infinitesimal Hyper-reals, or to the infinite Hyper-reals, or to the non-constant Hyper-reals.
16. No neighbourhood of a Hyper-real is homeomorphic to an \mathbb{R}^n ball. Therefore, the Hyper-real line is not a manifold.
17. The Hyper-real line is totally ordered like a line, but it is not spanned by one element, and it is not one-dimensional.

2.

Hyper-Complex Plane

Each complex number $\alpha + i\beta$ can be represented by a Cauchy sequence of rational complex numbers, $\langle r_1 + is_1, r_2 + is_2, r_3 + is_3, \dots \rangle$ so that $r_n + is_n \rightarrow \alpha + i\beta$.

The constant sequence $(\alpha + i\beta, \alpha + i\beta, \alpha + i\beta, \dots)$ is a Constant Hyper-Complex Number.

Following [Dan2] we claim that,

1. Any set of sequences $(l_1 + io_1, l_2 + io_2, l_3 + io_3, \dots)$, where (l_1, l_2, l_3, \dots) belongs to one family of infinitesimal hyper reals, and (o_1, o_2, o_3, \dots) belongs to another family of infinitesimal hyper-reals, constitutes a family of infinitesimal hyper-complex numbers.
2. Each hyper-complex infinitesimal has a polar representation $dz = (dr)e^{i\phi} = o_*e^{i\phi}$, where $dr = o_*$ is an infinitesimal, and $\phi = \arg(dz)$.
3. The infinitesimal hyper-complex numbers are smaller in length, than any complex number, yet strictly greater than

zero.

4. Their reciprocals $\left(\frac{1}{t_1+io_1}, \frac{1}{t_2+io_2}, \frac{1}{t_3+io_3}, \dots\right)$ are the infinite hyper-complex numbers.
5. The infinite hyper-complex numbers are greater in length than any complex number, yet strictly smaller than infinity.
6. The sum of a complex number with an infinitesimal hyper-complex is a non-constant hyper-complex.
7. The Hyper-Complex Numbers are the totality of constant hyper-complex numbers, a family of hyper-complex infinitesimals, a family of infinite hyper-complex, and non-constant hyper-complex.
8. The Hyper-Complex Plane is the direct product of a Hyper-Real Line by an imaginary Hyper-Real Line.
9. In Cartesian Coordinates, the Hyper-Real Line serves as an x coordinate line, and the imaginary as an iy coordinate line.
10. In Polar Coordinates, the Hyper-Real Line serves as a Range r line, and the imaginary as an $i\theta$ coordinate. Radial symmetry leads to Polar Coordinates.
11. The Hyper-Complex Plane includes the complex numbers separated by the non-constant hyper-complex

- numbers. Each complex number is the center of a disk of hyper-complex numbers, that includes no other complex number.
12. In particular, zero is separated from any complex number by a disk of complex infinitesimals.
 13. Zero is not a complex infinitesimal, because the length of zero is not strictly greater than zero.
 14. We do not add infinity to the hyper-complex plane.
 15. The hyper-complex plane is embedded in \mathbb{C}^∞ , and is not homeomorphic to the Complex Plane \mathbb{C} . There is no bi-continuous one-one mapping from the hyper-complex Plane onto the Complex Plane.
 16. In particular, there are no points in the Complex Plane that can be assigned uniquely to the hyper-complex infinitesimals, or to the infinite hyper-complex numbers, or to the non-constant hyper-complex numbers.
 17. No neighbourhood of a hyper-complex number is homeomorphic to a \mathbb{C}^n ball. Therefore, the Hyper-Complex Plane is not a manifold.
 18. The Hyper-Complex Plane is not spanned by two elements, and is not two-dimensional.

3.

Hyper-Complex Function

3.1 Definition of a hyper-complex function

$f(z)$ is a hyper-complex function, iff it is from the hyper-complex numbers into the hyper-complex numbers.

This means that any number in the domain, or in the range of a hyper-complex $f(x)$ is either one of the following

- complex
- complex + infinitesimal
- infinitesimal
- infinite hyper-complex

3.2 *Every function from complex numbers into complex numbers is a hyper-complex function.*

3.3 $\frac{\sin(dz)}{dz}$ *has the constant hyper-complex value 1*

Proof: $\sin(dz) = dz - \frac{(dz)^3}{3!} + \frac{(dz)^5}{5!} - \dots$

$$\frac{\sin(dz)}{dz} = 1 - \frac{(dz)^2}{3!} + \frac{(dz)^4}{5!} - \dots$$

3.4 $\cos(dz)$ has the constant hyper-complex value 1

Proof: $\cos(dz) = 1 - \frac{(dz)^2}{2!} + \frac{(dz)^4}{4!} - \dots$

3.5 e^{dz} has the constant hyper-complex value 1

Proof: $e^{dz} = 1 + dz + \frac{(dz)^2}{2!} + \frac{(dz)^3}{3!} + \frac{(dz)^4}{4!} + \dots$

3.6 $e^{\frac{1}{dz}}$ is an infinite hyper-complex, and $\left| e^{\frac{1}{dz}} \right| = e^{\frac{1}{dr} \cos \phi}$.

Proof: $\left| e^{\frac{1}{dz}} \right| = e^{\frac{1}{dr} \operatorname{Re}[e^{-i\phi}]} = e^{\frac{1}{dr} \cos \phi}$.

3.7 $\log(dz)$ is an infinite hyper-complex, and $|\log(dz)| > \frac{1}{dr}$

Proof: $|\log(dz)| = \sqrt{[\log(dr)]^2 + \phi^2} > \log(dr) > \frac{1}{dr}$

4.

Hyper-Complex Path Integral

Following the definition of the Hyper-real Integral in [Dan3], the Hyper-Complex Integral of $f(z)$ over a path $z(t)$, $t \in [\alpha, \beta]$, in its domain, is the sum of the areas $f(z)z'(t)dt = f(z)dz(t)$ of the rectangles with base $z'(t)dt = dz$, and height $f(z)$.

4.1 Hyper-Complex Path Integral Definition

Let $f(z)$ be hyper-complex function, defined on a domain in the Hyper-Complex Plane. The domain may not be bounded.

$f(z)$ may take infinite hyper-complex values, and need not be bounded.

Let $z(t)$, $t \in [\alpha, \beta]$, be a path, $\gamma(a, b)$, so that $dz = z'(t)dt$, and $z'(t)$ is continuous.

For each t , there is a hyper-complex rectangle with base $[z(t) - \frac{dz}{2}, z(t) + \frac{dz}{2}]$, height $f(z)$, and area $f(z(t))dz(t)$.

We form the **Integration Sum** of all the areas that start at $z(\alpha) = a$, and end at $z(\beta) = b$,

$$\sum_{t \in [\alpha, \beta]} f(z(t)) dz(t).$$

If for any infinitesimal $dz = z'(t)dt$, the Integration Sum equals the same hyper-complex number, then $f(z)$ is Hyper-Complex Integrable over the path $\gamma(a, b)$.

Then, we call the Integration Sum the Hyper-Complex Integral of $f(z)$ over the $\gamma(a, b)$, and denote it by $\int_{\gamma(a, b)} f(z) dz$.

If the hyper-complex number is an infinite hyper-complex, then it equals $\int_{\gamma(a, b)} f(z) dz$.

If the hyper-complex number is finite, then its constant part equals $\int_{\gamma(a, b)} f(z) dz$. \square

The Integration Sum may take infinite hyper-complex values, such as $\frac{1}{dz}$, but may not equal to ∞ .

The Hyper-Complex Integral of the function $f(z) = \frac{1}{|z|}$ over a path that goes through $z = 0$ diverges.

4.2 The Countability of the Integration Sum

In [Dan1], we established the equality of all positive infinities:

We proved that the number of the Natural Numbers,

$Card\mathbb{N}$, equals the number of Real Numbers, $Card\mathbb{R} = 2^{Card\mathbb{N}}$, and we have

$$Card\mathbb{N} = (Card\mathbb{N})^2 = \dots = 2^{Card\mathbb{N}} = 2^{2^{Card\mathbb{N}}} = \dots \equiv \infty.$$

In particular, we demonstrated that the real numbers may be well-ordered.

Consequently, there are countably many real numbers in the interval $[\alpha, \beta]$, and the Integration Sum has countably many terms.

While we do not sequence the real numbers in the interval, the summation takes place over countably many $f(z)dz$.

4.3 Continuous $f(z)$ is Path-Integrable

Hyper-Complex $f(z)$ Continuous on D is Path-Integrable on D

Proof:

Let $z(t)$, $t \in [\alpha, \beta]$, be a path, $\gamma(a, b)$, so that $dz = z'(t)dt$, and $z'(t)$ is continuous. Then,

$$f(z(t))z'(t) = (u(x(t), y(t)) + iv(x(t), y(t)))(x'(t) + iy'(t))$$

$$\begin{aligned}
&= \underbrace{\left[u(x(t), y(t))x'(t) - v(x(t), y(t))y'(t) \right]}_{U(t)} + \\
&\quad + i \underbrace{\left[u(x(t), y(t))y'(t) + v(x(t), y(t))x'(t) \right]}_{V(t)} \\
&= U(t) + iV(t),
\end{aligned}$$

where $U(t)$, and $V(t)$ are Hyper-Real Continuous on $[\alpha, \beta]$.

Therefore, by [Dan3, 12.4], $U(t)$, and $V(t)$ are integrable on $[\alpha, \beta]$.

Hence, $f(z(t))z'(t)$ is integrable on $[\alpha, \beta]$.

Since

$$\int_{t=\alpha}^{t=\beta} f(z(t))z'(t)dt = \int_{\gamma(a,b)} f(z)dz,$$

$f(z)$ is Path-Integrable on $\gamma(a, b)$. \square

5.

Hyper-real Delta Function

In [Dan5], we defined the Hyper-real Delta Function, and established its properties

1. The Delta Function is a Hyper-real function defined from the

Hyper-real line into the set of two Hyper-reals $\left\{0, \frac{1}{dx}\right\}$. The

Hyper-real 0 is the sequence $\langle 0, 0, 0, \dots \rangle$. The infinite Hyper-

real $\frac{1}{dx}$ depends on our choice of dx .

2. We will usually choose the family of infinitesimals that is

spanned by the sequences $\left\langle \frac{1}{n} \right\rangle, \left\langle \frac{1}{n^2} \right\rangle, \left\langle \frac{1}{n^3} \right\rangle, \dots$. It is a

semigroup with respect to vector addition, and includes all

the scalar multiples of the generating sequences that are

non-zero. That is, the family includes infinitesimals with

negative sign. Therefore, $\frac{1}{dx}$ will mean the sequence $\langle n \rangle$.

Alternatively, we may choose the family spanned by the

sequences $\left\langle \frac{1}{2^n} \right\rangle, \left\langle \frac{1}{3^n} \right\rangle, \left\langle \frac{1}{4^n} \right\rangle, \dots$. Then, $\frac{1}{dx}$ will mean the sequence $\left\langle 2^n \right\rangle$. Once we determined the basic infinitesimal dx , we will use it in the Infinite Riemann Sum that defines an Integral in Infinitesimal Calculus.

3. The Delta Function is strictly smaller than ∞

4. We define, $\delta(x) \equiv \frac{1}{dx} \chi_{\left[-\frac{dx}{2}, \frac{dx}{2}\right]}(x)$,

$$\text{where } \chi_{\left[-\frac{dx}{2}, \frac{dx}{2}\right]}(x) = \begin{cases} 1, & x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right] \\ 0, & \text{otherwise} \end{cases}.$$

5. Hence,

$$\diamond \text{ for } x < 0, \delta(x) = 0$$

$$\diamond \text{ at } x = -\frac{dx}{2}, \delta(x) \text{ jumps from } 0 \text{ to } \frac{1}{dx},$$

$$\diamond \text{ for } x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right], \delta(x) = \frac{1}{dx}.$$

$$\diamond \text{ at } x = 0, \delta(0) = \frac{1}{dx}$$

$$\diamond \text{ at } x = \frac{dx}{2}, \delta(x) \text{ drops from } \frac{1}{dx} \text{ to } 0.$$

$$\diamond \text{ for } x > 0, \delta(x) = 0.$$

$$\diamond x\delta(x) = 0$$

6. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \langle \chi_{[-\frac{1}{2}, \frac{1}{2}]}(x), 2\chi_{[-\frac{1}{4}, \frac{1}{4}]}(x), 3\chi_{[-\frac{1}{6}, \frac{1}{6}]}(x) \dots \rangle$

7. If $dx = \langle \frac{2}{n} \rangle$, $\delta(x) = \langle \frac{1}{2 \cosh^2 x}, \frac{2}{2 \cosh^2 2x}, \frac{3}{2 \cosh^2 3x}, \dots \rangle$

8. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \langle e^{-x} \chi_{[0, \infty)}, 2e^{-2x} \chi_{[0, \infty)}, 3e^{-3x} \chi_{[0, \infty)}, \dots \rangle$

9. $\int_{x=-\infty}^{x=\infty} \delta(x) dx = 1.$

6.

Hyper-complex Delta of a Complex Variable $\delta(z)$

In [Dan9], we introduced the Hyper-Complex Delta Function of a Complex Variable $\delta(z)$:

- 1) The Hyper-Complex Delta Function $\delta(z)$ is defined from the Hyper-Complex plane into the set of two hyper-complex numbers, $\left\{0, \frac{1}{2\pi i dz}\right\}$.

The hyper-complex 0 is the sequence $\langle 0, 0, 0, \dots \rangle$.

The infinite hyper-complex $\frac{1}{2\pi i} \frac{1}{dz} = \frac{1}{2\pi i} \frac{1}{dr} e^{-i\phi}$ depends on

$\text{Arg } z = \phi$. $\frac{1}{dr}$ will mean the sequence $\langle n \rangle$.

- 2) $\delta(z)$ is an infinite hyper-complex on the infinitesimal

hyper-complex disk $|z| \leq dr$. In particular, $\boxed{\delta(z) < \infty}$

$$\mathbf{3)} \quad \boxed{\delta(z - z_0) = \frac{1}{2\pi i} \frac{1}{dr} e^{-i \text{Arg}(z - z_0)} \chi_{\{|z - z_0| \leq dr\}}(z)},$$

$$\text{where } \chi_{\{|z - z_0| \leq dr\}}(z) = \begin{cases} 0, & |z - z_0| > dr \\ 1, & |z - z_0| \leq dr \end{cases}.$$

- on the disk, $|z - z_0| \leq dr$, $\delta(z - z_0) = \frac{1}{2\pi i} \frac{1}{dz}$.
- off the disk, for $|z - z_0| > dr$, $\delta(z - z_0) = 0$.

$$\mathbf{4)} \quad \boxed{(\delta(z))^n = \frac{1}{(2\pi i)^n} \frac{1}{(dr)^n} e^{-in\phi} \chi_{\{|z| \leq dr\}}(z)}, \quad n = 2, 3, \dots$$

$$\mathbf{5)} \quad \boxed{\delta(\zeta - z) = \frac{d}{dz} \frac{1}{2\pi i} (\text{Log}(\zeta - z)) \chi_{\{|\zeta - z| \leq dr\}}(\zeta)}$$

$$\mathbf{6)} \quad \boxed{\frac{d}{dz} \delta(\zeta - z) = \frac{1}{2\pi i} \frac{1}{(\zeta - z)^2} \chi_{\{|\zeta - z| \leq dr\}}(z)}$$

- in the disk $|\zeta - z| \leq dr$, $\frac{d}{dz} \delta(\zeta - z) = \frac{1}{2\pi i} \frac{1}{(dr)^2} e^{-2i\theta}$.
- off the disk, in $|\zeta - z| > dr$, $\frac{d}{dz} \delta(\zeta - z) = 0$.

$$\mathbf{7)} \quad \boxed{\frac{d^k}{dz^k} \delta(\zeta - z) = \frac{1}{2\pi i} \frac{k!}{(\zeta - z)^{k+1}} \chi_{\{|z| \leq dr\}}(z)}$$

- in the disk $|\zeta - z| \leq dr$, $\frac{d^k}{dz^k} \delta(\zeta - z) = \frac{k!}{2\pi i} \frac{1}{(dr)^{k+1}} e^{-i(k+1)\theta}$,
- off the disk, in $|\zeta - z| > dr$, $\frac{d^k}{dz^k} \delta(\zeta - z) = 0$.

$$\mathbf{8)} \quad \boxed{\delta(az) = \frac{1}{a} \delta(z)}$$

$$\mathbf{9)} \quad z_1 = \text{only zero of } f(z), \quad f'(z_1) \neq 0 \Rightarrow$$

$$\Rightarrow \boxed{\delta(f(z)) = \frac{1}{f'(z_1)} \delta(z - z_1)}$$

$$\mathbf{10)} \quad z_1, z_2 \text{ are the only zeros of } f(z); f'(z_1), f'(z_2) \neq 0 \Rightarrow$$

$$\Rightarrow \boxed{\delta(f(z)) = \frac{1}{f'(z_1)} \delta(z - z_1) + \frac{1}{f'(z_2)} \delta(z - z_2)}$$

$$\mathbf{11)} \quad \boxed{\delta(z^2 - a^2) = \frac{1}{2a} \delta(z - a) + \frac{1}{2a} \delta(z + a)}$$

$$\mathbf{12)} \quad \delta((z-a)(z-b)) = \frac{1}{a-b} \delta(z-a) + \frac{1}{b-a} \delta(z-b)$$

13) z_1, \dots, z_n are the only zeros of $f(z)$; $f'(z_1), \dots, f'(z_n) \neq 0 \Rightarrow$

$$\delta(f(z)) = \frac{1}{f'(z_1)} \delta(z - z_1) + \dots + \frac{1}{f'(z_n)} \delta(z - z_n)$$

14) z_1, z_2, \dots are zeros of $f(z)$, $f'(z_1), f'(z_2), \dots \neq 0 \Rightarrow$

$$\delta(f(z)) = \frac{1}{f'(z_1)} \delta(z - z_1) + \frac{1}{f'(z_n)} \delta(z - z_n) + \dots$$

15)

$$\delta(\sin z) = \dots + \delta(z + 2\pi) - \delta(z + \pi) + \delta(z) - \delta(z - \pi) + \delta(z - 2\pi) + \dots$$

$$\mathbf{16)} \quad \oint_{|\zeta-z|=dr} \delta(\zeta - z) d\zeta = 1$$

17) If $f(z)$ is Hyper-Complex Differentiable function at z

Then,
$$\oint_{|\zeta-z|=dr} f(\zeta) \delta(\zeta - z) d\zeta = f(z)$$

$$18) \quad \frac{d}{dz} f(z) = \oint_{|\zeta-z|=dr} f(\zeta) \frac{d}{dz} \delta(\zeta - z) d\zeta$$

$$19) \quad \frac{d^k}{dz^k} f(z) = \oint_{|\zeta-z|=dr} f(\zeta) \frac{d^k}{dz^k} \delta(\zeta - z) d\zeta$$

7.

Cauchy Integral Formula

7.1 Cauchy Integral Formula


If $f(z)$ is Hyper-Complex Differentiable function on a Hyper-Complex Simply-Connected Domain D .

Then,

$$f(z) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

for any loop γ , and any point z in its interior.

Proof: The Hyper-Complex function $\frac{f(\zeta)}{\zeta - z}$ is Differentiable on the Hyper-Complex Simply-Connected domain D , and on a path that includes γ and an infinitesimal circle about z .



Then, the integrals on the lines between γ and the circle have opposite signs and cancel each other.

The integral over the circle has a negative sign because its direction is clockwise, and by Cauchy Integral Theorem,

$$\oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta - \oint_{|\zeta-z|=dr} \frac{f(\zeta)}{\zeta - z} d\zeta = 0.$$

Therefore,

$$\begin{aligned} \oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta &= \oint_{|\zeta-z|=dr} \frac{f(\zeta)}{\zeta - z} d\zeta \\ &= 2\pi i \underbrace{\oint_{|\zeta-z|=dr} f(\zeta) \frac{1}{2\pi i} \frac{1}{\zeta - z} d\zeta}_{f(z)}. \square \end{aligned}$$

8.

Hyper-real Fourier Transform

In [Dan6], we defined the Fourier Transform and established its properties

1. $\mathcal{F}\{\delta(x)\} = 1$

2. $\delta(x) = \text{the inverse Fourier Transform of the unit function } 1$

$$= \frac{1}{2\pi} \int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega$$

$$= \int_{\nu=-\infty}^{\nu=\infty} e^{2\pi i x} d\nu, \quad \omega = 2\pi\nu$$

3. $\frac{1}{2\pi} \int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega \Big|_{x=0} = \frac{1}{dx} = \text{an infinite Hyper-real}$

$$\int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega \Big|_{x \neq 0} = 0$$

4. Fourier Integral Theorem

$$f(x) = \frac{1}{2\pi} \int_{k=-\infty}^{k=\infty} \left(\int_{\xi=-\infty}^{\xi=\infty} f(\xi) e^{-ik\xi} d\xi \right) e^{ikx} dk$$

does not hold in the Calculus of Limits, under any conditions.

5. Fourier Integral Theorem in Infinitesimal Calculus

If $f(x)$ is a Hyper-real function,

Then,

➤ *the Fourier Integral Theorem holds.*

$$\text{➤ } \int_{x=-\infty}^{x=\infty} f(x) e^{-i\alpha x} dx \text{ converges to } F(\alpha)$$

$$\text{➤ } \frac{1}{2\pi} \int_{\alpha=-\infty}^{\alpha=\infty} F(\alpha) e^{-i\alpha x} d\alpha \text{ converges to } f(x)$$

6. 2-Dimesional Fourier Transform

$$\begin{aligned} \mathcal{F} \{ f(x, y) \} &= \int_{y=-\infty}^{y=\infty} \int_{x=-\infty}^{x=\infty} f(x, y) e^{-i\omega_x x - i\omega_y y} dx dy \\ &= \int_{y=-\infty}^{y=\infty} \int_{x=-\infty}^{x=\infty} f(x, y) e^{-2\pi i(\nu_x x + \nu_y y)} dx dy, \quad \begin{aligned} \omega_x &= 2\pi\nu_x \\ \omega_y &= 2\pi\nu_y \end{aligned} \end{aligned}$$

7. 2-Dimesional Inverse Fourier Transform

$$\begin{aligned}\mathcal{F}^{-1}\{F(\omega_x, \omega_y)\} &= \frac{1}{(2\pi)^2} \int_{\omega_y=-\infty}^{\omega_y=\infty} \int_{\omega_x=-\infty}^{\omega_x=\infty} F(\omega_x, \omega_y) e^{i(\omega_x x + \omega_y y)} d\omega_x d\omega_y \\ &= \int_{\nu_y=-\infty}^{\nu_y=\infty} \int_{\nu_x=-\infty}^{\nu_x=\infty} F(2\pi\nu_x, 2\pi\nu_y) e^{2\pi i(\nu_x x + \nu_y y)} d\nu_x d\nu_y, \quad \begin{aligned} \omega_x &= 2\pi\nu_x \\ \omega_y &= 2\pi\nu_y \end{aligned}\end{aligned}$$

8. 2-Dimesional Fourier Integral Theorem

$$\begin{aligned}f(x, y) &= \frac{1}{(2\pi)^2} \int_{\omega_y=-\infty}^{\omega_y=\infty} \int_{\omega_x=-\infty}^{\omega_x=\infty} \left(\int_{\eta=-\infty}^{\eta=\infty} \int_{\xi=-\infty}^{\xi=\infty} f(\xi, \eta) e^{-i\omega_x \xi - i\omega_y \eta} d\xi d\eta \right) e^{i(\omega_x x + \omega_y y)} d\omega_x d\omega_y \\ &= \int_{\eta=-\infty}^{\eta=\infty} \int_{\xi=-\infty}^{\xi=\infty} f(\xi, \eta) \left(\frac{1}{2\pi} \int_{\omega_x=-\infty}^{\omega_x=\infty} e^{i\omega_x(x-\xi)} d\omega_x \right) d\xi \left(\frac{1}{2\pi} \int_{\omega_y=-\infty}^{\omega_y=\infty} e^{i\omega_y(y-\eta)} d\omega_y \right) d\eta \\ &= \int_{\eta=-\infty}^{\eta=\infty} \int_{\xi=-\infty}^{\xi=\infty} f(\xi, \eta) \left(\int_{\nu_x=-\infty}^{\nu_x=\infty} e^{2\pi i\nu_x(x-\xi)} d\nu_x \right) d\xi \left(\int_{\nu_y=-\infty}^{\nu_y=\infty} e^{2\pi i\nu_y(y-\eta)} d\nu_y \right) d\eta, \quad \begin{aligned} \omega_x &= 2\pi\nu_x \\ \omega_y &= 2\pi\nu_y \end{aligned}\end{aligned}$$

9. 2-Dimesional Delta Function


$$\begin{aligned}\delta(x, y) &= \left(\frac{1}{2\pi} \int_{\omega_x=-\infty}^{\omega_x=\infty} e^{i\omega_x x} d\omega_x \right) \left(\frac{1}{2\pi} \int_{\omega_y=-\infty}^{\omega_y=\infty} e^{i\omega_y y} d\omega_y \right) \\ &= \left(\int_{\nu_x=-\infty}^{\nu_x=\infty} e^{2\pi i\nu_x x} d\nu_x \right) \left(\int_{\nu_y=-\infty}^{\nu_y=\infty} e^{2\pi i\nu_y y} d\nu_y \right), \quad \begin{aligned} \omega_x &= 2\pi\nu_x \\ \omega_y &= 2\pi\nu_y \end{aligned}\end{aligned}$$

9.

Fourier Transform of a Hyper-Complex $f(z)$

For a fixed hyper-complex ω , and a varying hyper-real θ , the exponential $e^{-i\omega\theta}$ spirals along its multi-plane Riemann Surface. Such surface can be graphed by say

```
ParametricPlot3D[{3 Cos[t], 3 Sin[t], 1/11 t}, {t, -10 π, 10 π}]
```



For a hyper-complex function $f(\theta) = u(\theta) + iv(\theta)$, the infinitesimal products $f(\theta)e^{-i\omega\theta} d\theta$, sum up to the Fourier Transform of $f(\theta)$, the

Integration Sum

$$\sum_{\theta=-\infty}^{\theta=\infty} f(\theta)e^{-i\omega\theta}d\theta.$$

When θ is replaced with a varying hyper-complex z , $-\infty$, and ∞ , coincide into one infinity denoted by ∞ .

Then, an integration path from $-\infty$, to ∞ is a closed path through ∞ , which, by Cauchy Theorem, may be homologous to a circle in the Complex plane.

In the Calculus of Limits, it is erroneously believed [Dan4] that the Fourier Transform requires boundedness conditions on the function.

Extending these unnecessary conditions to the Complex plane, Titchmarsh [Titchmarsh, p.42] defines the Fourier Transform of $f(z)$ along a path parallel to the x axis.

Then, the Fourier Transform is an analytic function in a strip parallel to the x axis. And at the borders of the strip, the Analytic Function which is inseparable from its disk of convergence, will have one sided derivative.

No one, including Titchmarsh, cared to explore this unnatural geometry, and the Fourier Transform that is analytic in a strip parallel to the x axis, was never developed further.

We define the Fourier Transform of a hyper-complex function $f(z)$ along a closed path γ by the Integration Sum

$$\sum_{z \in \gamma} f(z) e^{-i\omega z} dz.$$

We will see that The existence of the Fourier Transform of an Analytic Function follows from the existence of the Fourier Integral of that Function.

We will establish that The Fourier Integral of an Analytic $f(z)$ is its Cauchy Integral Formula.

In that Formula, the function's values are swept by the Hyper-Complex Delta Function, as in Section 6.17.

To obtain the Fourier Integral of an Analytic Function, we need the Fourier Integral of the Complex Delta Function.

To that end, we proceed with the Fourier Transform of the Complex Delta function.

10.

Fourier Transform of $\delta(z)$

10.1 Fourier Transform of $\delta(z)$

$$\mathcal{F}\{\delta(z)\} = \sum_{z \in \gamma} \delta(z) e^{-i\omega z} dz.$$

Without loss of generality, γ may be the unit circle $z = e^{i\phi}$.

Therefore,

$$\mathbf{10.2} \quad \boxed{\mathcal{F}\{\delta(z)\} = 1}$$

Proof:
$$\mathcal{F}\{\delta(z)\} = \oint_{|z|=1} \delta(z) e^{-i\omega z} dz$$

By Cauchy Theorem,

$$= \frac{1}{2\pi i} \oint_{|z|=dr} \frac{1}{z} e^{-i\omega z} dz$$

By Cauchy Integral Theorem,

$$= e^{-i\omega z} \Big|_{z=0}$$

$$= 1. \square$$

11.

Fourier Integral of $\delta(z)$

Since the complex Delta is the inverse Transform of $\hat{f}(\omega) = 1$, we have

11.1 Fourier Integral of $\delta(z)$

$$\delta(z) = \frac{1}{2\pi} \oint_{|\omega|=\eta} e^{i\omega z} d\omega,$$

where η is an infinitesimal

By Cauchy Integral Theorem, the integration path may be along the unit circle.

Hence,

11.2 Fourier Integral of $\delta(z - \zeta)$

$$\delta(\zeta - z) = \frac{1}{2\pi} \oint_{|\omega|=1} e^{i\omega(\zeta-z)} d\omega$$

12.

Fourier Integral of an Analytic $f(z)$

12.1 Fourier Integral Theorem for hyper-complex Analytic $f(z)$, along infinitesimal paths

If $f(z)$ is a hyper-complex analytic function,

Then, the Complex Fourier Integral Theorem holds:

$$f(z) = \frac{1}{2\pi} \oint_{|\omega|=\eta} \left(\oint_{|\zeta-z|=\varepsilon} f(\zeta) e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega,$$

where ε , and η are infinitesimals

Proof:

By the Cauchy Integral Formula, 5.1,

$$f(z) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

By Cauchy Integral Theorem,

$$= \frac{1}{2\pi i} \oint_{|\zeta-z|=dr} \frac{f(\zeta)}{\zeta - z} d\zeta,$$

By the definition of $\delta(z)$,

$$= \oint_{|\zeta-z|=\varepsilon} f(\zeta)\delta(z-\zeta)d\zeta,$$

Substituting from 11.2, $\delta(\zeta-z) = \frac{1}{2\pi} \oint_{|\omega|=dr} e^{i\omega(\zeta-z)}d\omega$, we have

$$f(z) = \frac{1}{2\pi} \oint_{|\zeta-z|=\varepsilon} f(\zeta) \left(\oint_{|\omega|=\eta} e^{i\omega z} d\omega \right) e^{-i\omega\zeta} d\zeta.$$

By changing the Summation order,

$$f(z) = \frac{1}{2\pi} \oint_{|\omega|=\eta} \left(\oint_{|\zeta-z|=\varepsilon} f(\zeta)e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega. \square$$

$$= \frac{1}{2\pi} \oint_{|\omega|=1} \left(\oint_{|\zeta-z|=1} f(\zeta)e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega$$

Similarly, we obtain

12.2 Fourier Integral Theorem for $f(z)$ on Unit Circles

If $f(z)$ is a hyper-complex analytic function, in a hyper-complex

domain that contains the unit circle $|\zeta-z|=1$

Then, the Complex Fourier Integral Theorem holds.

$$f(z) = \frac{1}{2\pi} \oint_{|\omega|=1} \left(\oint_{|\zeta-z|=1} f(\zeta)e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega$$

Proof: As above,

$$f(z) = \oint_{|\zeta-z|=\varepsilon} f(\zeta)\delta(z-\zeta)d\zeta.$$

Substituting from 11.2, $\delta(\zeta-z) = \frac{1}{2\pi} \oint_{|\omega|=1} e^{i\omega(\zeta-z)}d\omega$, we have

$$f(z) = \frac{1}{2\pi} \oint_{|\zeta-z|=\varepsilon} f(\zeta) \left(\oint_{|\omega|=1} e^{i\omega z} d\omega \right) e^{-i\omega\zeta} d\zeta.$$

By changing the Summation order,

$$f(z) = \frac{1}{2\pi} \oint_{|\omega|=1} \left(\oint_{|\zeta-z|=\varepsilon} f(\zeta)e^{-i\omega\zeta} d\zeta \right) e^{i\omega z} d\omega.$$

By the Cauchy Theorem, the integration along the infinitesimal circle, can be done along the unit circle. Hence

$$f(z) = \frac{1}{2\pi} \oint_{|\omega|=1} \left(\oint_{|\zeta-z|=1} f(\zeta)e^{-i\omega\zeta} d\zeta \right) e^{i\omega z} d\omega. \square$$

It follows that

12.3 For an Analytic Hyper-Complex Function $f(z)$, the

Fourier Integral is the Cauchy Integral Formula for $f(z)$

13.

Fourier Transform of Analytic $f(z)$

The convergence of the Fourier Integral representing a hyper-complex analytic function $f(z)$, implies the existence of the Fourier Transform of $f(z)$, and its inverse transform

13.1

If $f(z)$ is a hyper-complex analytic function on a hyper-complex domain that includes the circle $|\zeta - z| = 1$,

Then,

1) the hyper-complex integral

$$\oint_{|\zeta-z|=1} f(\zeta)e^{-i\omega\zeta}d\zeta$$

converges to $\hat{f}(\omega)$

2) the hyper-complex integral

$$\frac{1}{2\pi} \oint_{|\omega|=1} \hat{f}(\omega)e^{iz\omega}d\omega$$

converges to $f(z)$

Proof:

The convergence

$$\frac{1}{2\pi} \oint_{|\omega|=1} \left(\oint_{|\zeta-z|=\varepsilon} f(\zeta) e^{-i\omega\zeta} d\zeta \right) e^{iz\omega} d\omega = f(z)$$

mandates that

1) The Complex Fourier Transform of $f(z)$,

$$\oint_{|\zeta-z|=\varepsilon} f(\zeta) e^{-i\omega\zeta} d\zeta,$$

converges to a hyper-complex function $\hat{f}(\omega)$, some of its values may be infinite hyper-complex, like the complex Delta Function.

2) The Inverse Fourier Transform of the main branch of $\hat{f}(\omega)$

$$\frac{1}{2\pi} \oint_{|\omega|=\eta} \hat{f}(\omega) e^{iz\omega} d\omega$$

converges to the hyper-complex function $f(z)$. \square

References

[Bremermann] Hans Bremermann, “*Distributions, Complex Variables, and Fourier Transforms*” Addison-Wesley, 1965

[Dan1] Dannon, H. Vic, “[*Well-Ordering of the Reals, Equality of all Infinities, and the Continuum Hypothesis*](#)” in Gauge Institute Journal Vol.6 No 2, May 2010;

[Dan2] Dannon, H. Vic, “[*Infinitesimals*](#)” in Gauge Institute Journal Vol.6 No 4, November 2010;

[Dan3] Dannon, H. Vic, “[*Infinitesimal Calculus*](#)” in Gauge Institute Journal Vol.7 No 4, November 2011;

[Dan4] Dannon, H. Vic, “[*The Delta Function*](#)” in Gauge Institute Journal Vol.8 No 1, February 2012;

[Dan5] Dannon, H. Vic, “[*Infinitesimal Vector Calculus*](#)” in Gauge Institute Journal

[Dan6] Dannon, H. Vic, “[*Circular and Spherical Delta Functions*](#)” in Gauge Institute Journal

[Dan7] Dannon, H. Vic, “[*Infinitesimal Complex Calculus*](#)” in Gauge Institute Journal, Vol. 10 No. 4, November 2014.

[Dan8] H. Vic Dannon, “[*Delta Function, the Fourier Transform, and Fourier Integral Theorem*](#)”, in Gauge Institute Journal Vol.8 No 2, May 2012.

[Dan9] H. Vic Dannon, “[*Delta Function of a Complex Variable*](#)” in Gauge Institute Journal Vol.11 No 1, February 2015.

[Needham] Tristan Needham, “*Visual Complex Analysis*” Oxford U. Press, 1998 (with corrections)

[Paley, Wiener] Raymond Paley, and Norbert Wiener, “*Fourier Transforms in the Complex Plane*” American Mathematical Society, 1934

[Sneddon] Ian Sneddon, “*Fourier Transforms*”, McGraw-Hill, 1959.

[Titchmarsh] E. C. Titchmarsh “*Introduction to the theory of Fourier Integrals*”, Third Edition, Chelsea, 1986.