

Delta Function, and Expansion in Laguerre Functions

H. Vic Dannon
vic0@comcast.net
June, 2012

Abstract Let $f(x)$ be defined on $[0, \infty)$, and let $L_n(x)$ be the Laguerre Polynomials on $[0, \infty)$,

$$L_0(x) = 1, L_1(x) = -x + 1, L_2(x) = \frac{1}{2}x^2 - 2x + 1,$$

$$L_3(x) = -\frac{1}{6}x^3 + \frac{3}{2}x^2 - 3x + 1, \dots$$

The Laguerre Series associated with $f(x)$ is

$$a_0L_0(x) + a_1L_1(x) + a_2L_2(x) + \dots$$

where

$$a_n = \int_{\xi=-\infty}^{\xi=\infty} e^{-x} f(\xi) L_n(\xi) d\xi$$

are the Laguerre coefficients.

The Laguerre Series Theorem supplies the conditions under which the Laguerre Series associated with $f(x)$ equals $f(x)$.

It is believed to hold in the Calculus of Limits for smooth enough function. In fact,

*The Theorem cannot be proved in the Calculus of Limits
under any conditions,*

because the summation of the Laguerre Series requires integration of the singular Laguerre Kernel.

Plots of partial sums of the Laguerre Series speak volumes about the sensibility of the claims to have infinity bound by epsilon.

In Infinitesimal Calculus, the Laguerre Kernel

$$e^{-x} \{ L_0(\xi)L_0(x) + L_1(\xi)L_1(x) + L_2(\xi)L_2(x) + \dots \}$$

is the Delta Function $\delta(\xi - x)$.

$\delta(\xi - x)$ equals its Laguerre Series, and the Laguerre Series associated with any hyper-real integrable $f(x)$, equals $f(x)$

Keywords: Infinitesimal, Infinite-Hyper-Real, Hyper-Real, infinite Hyper-real, Infinitesimal Calculus, Delta Function, Laguerre Polynomials, Laguerre Coefficients, Delta Function, Laguerre Series, Laguerre Kernel, Expansion in Laguerre Functions,

2000 Mathematics Subject Classification 26E35; 26E30;
26E15; 26E20; 26A06; 26A12; 03E10; 03E55; 03E17; 03H15;
46S20; 97I40; 97I30.

Contents

0. The Origin of the Laguerre Series Theorem
1. Divergence of the Laguerre Kernel in the Calculus of Limits
2. Hyper-real line.
3. Integral of a Hyper-real Function
4. Delta Function
5. Convergent Series
6. Laguerre Sequence and $\delta(\xi - x)$
7. Laguerre Kernel and $\delta(\xi - x)$.
8. Laguerre Series of $\delta(\xi - x)$
9. Laguerre Series Theorem

References

The Origin of the Laguerre Series

Theorem

The Laguerre Polynomials on $[0, \infty)$

$$L_0(x) = 1, L_1(x) = -x + 1, L_2(x) = \frac{1}{2}x^2 - 2x + 1,$$

$$L_3(x) = -\frac{1}{6}x^3 + \frac{3}{2}x^2 - 3x + 1, \dots$$

are orthogonal so that

$$\int_{x=0}^{x=\infty} e^{-x} L_m(x) L_n(x) dx = \delta_{mn}.$$

0.1 Laguerre

generated the polynomials

$$\tilde{L}_n(x) = n! L_n(x),$$

by expanding

$$\frac{1}{1-\alpha} e^{-\frac{\alpha}{1-\alpha}x} = \tilde{L}_0(x) + \frac{1}{1!} \tilde{L}_1(x)\alpha + \frac{1}{2!} \tilde{L}_2(x)\alpha^2 + \frac{1}{3!} \tilde{L}_3(x)\alpha^3 + \dots,$$

with

$$|\alpha| < 1. \text{ Hence, } \frac{1}{1-\alpha} = 1 + \alpha + \alpha^2 + \alpha^3 + \dots$$

We expand

$$\frac{1}{1-\alpha} e^{-\frac{\alpha}{1-\alpha}x} = L_0(x) + L_1(x)\alpha + L_2(x)\alpha^2 + L_3(x)\alpha^3 + \dots$$

and equate it to

$$\frac{1}{1-\alpha} e^{-\frac{\alpha}{1-\alpha}x} = \frac{1}{\underbrace{1-\alpha}_{1+\alpha+\alpha^2+\alpha^3+\dots}} - \frac{\alpha}{\underbrace{(1-\alpha)^2}_{\alpha(1+\alpha+\alpha^2+\dots)^2}} x + \frac{1}{2!} \frac{\alpha^2}{\underbrace{(1-\alpha)^3}_{\alpha^2(1+\alpha+\dots)^3}} x^2 - \frac{1}{3!} \frac{\alpha^3}{\underbrace{(1-\alpha)^4}_{\alpha^3(1+\alpha+\dots)^4}} x^3 \dots$$

$$\begin{aligned}
&= 1 + \alpha + \alpha^2 + \alpha^3 + \dots - \alpha \frac{(1 + \alpha + \alpha^2 + \dots)^2}{1 + \alpha^2 + \alpha^4 + 2\alpha + 2\alpha^2 + 2\alpha^3} x + \frac{1}{2!} \alpha^2 \frac{(1 + \alpha + \dots)^3}{1 + 3\alpha + 3\alpha^2 + \alpha^3 + \dots} x^2 + \\
&\qquad\qquad\qquad - \frac{1}{3!} \alpha^2 \frac{(1 + \alpha + \dots)^3}{1 + 3\alpha + 3\alpha^2 + \alpha^3 + \dots} x^2 + \dots \\
&= \underbrace{1}_{L_0(x)} + \underbrace{(1 - x)\alpha}_{L_1(x)} + \underbrace{(1 - 2x + \frac{1}{2}x^2)\alpha^2}_{L_2(x)} + \underbrace{(1 - 3x + \frac{3}{2}x^2 - \frac{1}{6}x^3)\alpha^3}_{L_3(x)} + \dots
\end{aligned}$$

0.2 Schrodinger Equation for an electron orbiting a proton

An electron with mass m , orbits a proton, at a distance r , under the force $k \frac{e^2}{r^2}$.

The potential energy of the electron is

$$V(r) = -k \frac{e^2}{r}, \quad \text{where } k = \frac{1}{4\pi\epsilon_0}$$

in the Meter-Kilogram-Second-Volt-Ampere system.

De Broglie associated with the electron a wave of length

$$\lambda = \frac{h}{mv},$$

where v is the velocity of the electron, and h is Planck's constant.

The wave's frequency is

$$\nu = \frac{v}{\lambda} = \frac{v}{\frac{h}{mv}} = \frac{mv^2}{h}$$

The wave's angular frequency is

$$\omega = 2\pi\nu = 2\pi \frac{mv^2}{h}$$

In terms of the De Broglie wave, the electron's energy is a multiple of Planck's radiation energy. Being binding energy, it is taken with a negative sign. That is,

$$E = -\varepsilon h\nu = -\varepsilon \hbar\omega, \quad \hbar = \frac{h}{2\pi}, \quad \varepsilon \text{ is the multiplier.}$$

The kinetic energy of the electron is

$$\frac{1}{2}mv^2 = E - V.$$

Hence,

$$mv = \sqrt{2m(E - V)},$$

$$\lambda = \frac{h}{\sqrt{2m(E - V)}},$$

$$v = \lambda \frac{\nu}{\frac{1}{2\pi}\omega} = \frac{\hbar\omega}{\sqrt{2m(E - V)}}.$$

$$\frac{1}{v^2} = \frac{2m(E - V)}{\hbar^2\omega^2}$$

Schrodinger postulated a complex valued potential

$$\Psi(r, \theta, \phi, t) = \psi(r, \theta, \phi)e^{i\omega t}$$

that satisfies the wave equation

$$\nabla^2\Psi(r, \theta, \phi, t) = \frac{1}{v^2}\partial_t^2\Psi(r, \theta, \phi, t).$$

Then,

$$\begin{aligned}
0 &= \nabla^2 \Psi(r, \theta, \phi, t) - \frac{1}{v^2} \partial_t^2 \Psi(r, \theta, \phi, t) \\
&= \nabla^2 \psi(r, \theta, \phi) e^{i\omega t} - \frac{2m(E - V)}{\hbar^2 \omega^2} \psi(r, \theta, \phi) (-\omega^2) e^{i\omega t}.
\end{aligned}$$

The Schrodinger equation for the orbiting electron is

$$\nabla^2 \psi(r, \theta, \phi) + \frac{2m}{\hbar^2} (E - V) \psi(r, \theta, \phi) = 0.$$

Substituting E , and V ,

$$\nabla^2 \psi(r, \theta, \phi) + \frac{2m}{\hbar^2} (-\varepsilon \hbar \omega + k \frac{e^2}{r}) \psi(r, \theta, \phi) = 0$$

In radial coordinates

$$\frac{1}{r^2} \partial_r (r^2 \partial_r \psi) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \psi) + \frac{1}{r^2 \sin^2 \theta} \partial_\phi^2 \psi + \frac{2m}{\hbar^2} (\varepsilon \hbar \omega + k \frac{e^2}{r}) \psi = 0.$$

Assuming that

$$\psi(r, \theta, \phi) = R(r) \Theta(\theta) \Phi(\phi),$$

the Schrodinger equation becomes

$$\frac{1}{R} (r^2 R')' + \frac{1}{\Theta \sin \theta} (\sin \theta \Theta')' + \frac{1}{\Phi \sin^2 \theta} \Phi'' + \frac{2m}{\hbar^2} r^2 (-\varepsilon \hbar \omega + k \frac{e^2}{r}) = 0.$$

Then,

$$\frac{1}{R} (r^2 R')' + \frac{2m}{\hbar^2} r^2 (-\varepsilon \hbar \omega + k \frac{e^2}{r}) = \text{const} \equiv C_1.$$

Substituting

$$R(r) = r^l,$$

$$l(l+1) + \frac{2m}{\hbar^2} r^2 (-\varepsilon \hbar \omega + k \frac{e^2}{r}) = C_1.$$

Hence,

$$l(l+1) = C_1.$$

Thus, the Schrodinger Radial Equation is

$$\begin{aligned} \frac{1}{R}(r^2 R')' + \frac{2m}{\hbar^2} r^2 (-\varepsilon \hbar \omega + k \frac{e^2}{r}) &= l(l+1) \\ r^2 R'' + 2r R' + \frac{2m}{\hbar^2} r^2 (-\varepsilon \hbar \omega + k \frac{e^2}{r}) R &= l(l+1)R, \\ r^2 R'' + 2r R' + \frac{2m}{\hbar^2} r^2 [-\varepsilon \hbar \omega + k \frac{e^2}{r} - \frac{\hbar^2}{2mr^2} l(l+1)] R &= 0 \\ R'' + \frac{2}{r} R' + \frac{2m}{\hbar^2} [-\varepsilon \hbar \omega + k \frac{e^2}{r} - \frac{\hbar^2}{2m} \frac{1}{r^2} l(l+1)] R &= 0. \end{aligned}$$

Multiplying by $\frac{\hbar}{8m\omega}$,

$$\begin{aligned} \frac{\hbar}{8m\omega} R'' + \frac{2}{r} \frac{\hbar}{8m\omega} R' + \frac{2m}{\hbar^2} \frac{\hbar}{8m\omega} [-\varepsilon \hbar \omega + \frac{1}{r} k e^2 - \frac{\hbar^2}{2m} \frac{1}{r^2} l(l+1)] R &= 0, \\ \underbrace{\frac{\hbar}{8m\omega} R''(r)}_{R''(\rho)} + \underbrace{\frac{2}{r} \frac{\hbar}{8m\omega} R'(r)}_{\frac{2}{\rho} R'(\rho)} + \\ + [-\frac{1}{4} \varepsilon + \frac{1}{r} \underbrace{\sqrt{\frac{\hbar}{8m\omega}}}_{\frac{1}{\rho}} \underbrace{\sqrt{\frac{\hbar}{8m\omega} \frac{2m}{\hbar^2} k e^2}}_{\sqrt{\frac{m}{2\hbar\omega} \frac{k e^2}{\hbar}}} - \frac{1}{r^2} \underbrace{\frac{\hbar}{8m\omega}}_{\frac{1}{\rho^2}} l(l+1)] R &= 0 \end{aligned}$$

The change of variable $\rho = \sqrt{\frac{8m\omega}{\hbar}} r$, gives

$$\frac{2m}{\hbar^2} \frac{\hbar}{8m\omega} \frac{1}{r} k e^2 = \frac{1}{\underbrace{r}_{\frac{1}{\rho}}} \underbrace{\sqrt{\frac{\hbar}{8m\omega}}}_{\frac{1}{\rho}} \underbrace{\sqrt{\frac{\hbar}{8m\omega} \frac{2m}{\hbar^2} k e^2}}_{\sqrt{\frac{m}{2\hbar\omega} \frac{k e^2}{\hbar}}} = \frac{1}{\rho} \sqrt{\frac{m}{2\hbar\omega} \frac{k e^2}{\hbar}}$$

$$R'(r) = \frac{dR}{d\rho} \frac{d\rho}{dr} = R'(\rho) \sqrt{\frac{8m\omega}{\hbar}},$$

$$\frac{2}{r} \frac{\hbar}{8m\omega} R'(r) = 2 \frac{1}{r} \underbrace{\sqrt{\frac{\hbar}{8m\omega}}}_{\frac{1}{\rho}} \underbrace{\sqrt{\frac{\hbar}{8m\omega}} R'(r)}_{R'(\rho)}$$

$$R''(r) = \frac{d}{d\rho} \left\{ R'(\rho) \sqrt{\frac{8m\omega}{\hbar}} \right\} \frac{d\rho}{dr} = R''(\rho) \frac{8m\omega}{\hbar},$$

and the equation becomes

$$R''(\rho) + \frac{2}{\rho} R'(\rho) + \left\{ -\frac{1}{4} \varepsilon + \frac{1}{\rho} \sqrt{\frac{m}{2\hbar\omega}} \frac{ke^2}{\hbar} - \frac{1}{\rho^2} l(l+1) \right\} R = 0.$$

For $\rho \rightarrow \infty$, the Radial Schrodinger equation

$$R''(\rho) + \frac{2}{\rho} R'(\rho) + \left\{ -\frac{1}{4} \varepsilon + \frac{1}{\rho} \sqrt{\frac{m}{2\hbar\omega}} \frac{ke^2}{\hbar} - \frac{1}{\rho^2} l(l+1) \right\} R = 0$$

becomes

$$R''(\rho) - \frac{1}{4} \varepsilon R = 0$$

Factoring

$$(D_\rho - \frac{1}{2} \sqrt{\varepsilon})(D_\rho + \frac{1}{2} \sqrt{\varepsilon})R(\rho) = 0.$$

we solve

$$(D_\rho - \frac{1}{2} \sqrt{\varepsilon})R(\rho) = 0$$

$$\frac{R'}{R} = \frac{1}{2} \sqrt{\varepsilon}$$

$$\log R = \frac{1}{2} \sqrt{\varepsilon} \rho + c$$

$$R_1 = C e^{\frac{1}{2} \sqrt{\varepsilon} \rho}.$$

As $\rho \rightarrow \infty$, $R_1 \rightarrow \infty$, and is discarded.

$$(D_\rho + \frac{1}{2}\sqrt{\varepsilon})R(\rho) = 0$$

$$\frac{R'}{R} = -\frac{1}{2}\sqrt{\varepsilon}$$

$$\log R = -\frac{1}{2}\sqrt{\varepsilon}\rho + c$$

$$R_2 = Ce^{-\frac{1}{2}\sqrt{\varepsilon}\rho}.$$

0.3 Laguerre Differential Equation

Denote

$$\sqrt{\frac{m}{2\hbar\omega}} \frac{ke^2}{\hbar} \equiv \lambda,$$

and substitute

$$R(\rho) = L(\rho)\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho}$$

in

$$R''(\rho) + \frac{2}{\rho}R'(\rho) + [-\frac{1}{4}\varepsilon + \frac{1}{\rho}\underbrace{\sqrt{\frac{m}{2\hbar\omega}} \frac{ke^2}{\hbar}}_\lambda - \frac{1}{\rho^2}l(l+1)]R = 0.$$

Then,

$$\begin{aligned} 0 &= (L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho})'' + \frac{2}{\rho}(L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho})' + [-\frac{1}{4}\varepsilon + \frac{1}{\rho}\lambda - \frac{1}{\rho^2}l(l+1)]L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} \\ &= (L'\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + Ll\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho})' \\ &\quad + \frac{2}{\rho}(L'\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + Ll\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho}) \\ &\quad + [-\frac{1}{4}\varepsilon + \frac{1}{\rho}\lambda - \frac{1}{\rho^2}l(l+1)]L\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} \\ &= L''\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + L'\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}L'\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + \end{aligned}$$

$$\begin{aligned}
& +L'l\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + Ll(l-1)\rho^{l-2}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}Ll\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} \\
& -\frac{1}{2}\sqrt{\varepsilon}L'\rho^le^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}Ll\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + \frac{1}{4}\varepsilon L\rho^le^{-\frac{1}{2}\sqrt{\varepsilon}\rho} \\
& +\frac{2}{\rho}(L'\rho^le^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + Ll\rho^{l-1}e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} - \frac{1}{2}\sqrt{\varepsilon}L\rho^le^{-\frac{1}{2}\sqrt{\varepsilon}\rho}) \\
& +[-\frac{1}{4}\varepsilon + \frac{1}{\rho}\lambda - \frac{1}{\rho^2}l(l+1)]L\rho^le^{-\frac{1}{2}\sqrt{\varepsilon}\rho}.
\end{aligned}$$

$$\begin{aligned}
0 & = L''\rho^l + L'l\rho^{l-1} - \frac{1}{2}\sqrt{\varepsilon}L'\rho^l + \\
& +L'l\rho^{l-1} + Ll(l-1)\rho^{l-2} - \frac{1}{2}\sqrt{\varepsilon}Ll\rho^{l-1} \\
& -\frac{1}{2}\sqrt{\varepsilon}L'\rho^l - \frac{1}{2}\sqrt{\varepsilon}Ll\rho^{l-1} + \frac{1}{4}\varepsilon L\rho^l \\
& +2L'\rho^{l-1} + 2Ll\rho^{l-2} - \sqrt{\varepsilon}L\rho^{l-1} \\
& -\frac{1}{4}\varepsilon L\rho^l + \lambda L\rho^{l-1} - l(l+1)L\rho^{l-2}.
\end{aligned}$$

The Schrodinger equation becomes Laguerre Differential Equation

$$\rho L'' + [2(l+1) - \sqrt{\varepsilon}\rho]L' + [\lambda - \sqrt{\varepsilon}(l+1)]L = 0,$$

Substituting in it

$$L(\rho) = c_0 + c_1\rho + c_2\rho^2 + \dots + c_j\rho^j + c_{j+1}\rho^{j+1} + c_{j+2}\rho^{j+2} + \dots,$$

we have

$$\rho \underbrace{D_\rho^2 \sum_{j=0}^{j=\infty} c_j \rho^j}_{\sum_{j=2}^{j=\infty} (j-1)j c_j \rho^{j-2}} + [2(l+1) - \sqrt{\varepsilon}\rho] \underbrace{D_\rho \sum_{j=0}^{j=\infty} c_j \rho^j}_{\sum_{j=1}^{j=\infty} j c_j \rho^{j-1}} + [\lambda - \sqrt{\varepsilon}(l+1)] \sum_{j=0}^{j=\infty} c_j \rho^j = 0,$$

$$\sum_{j=0}^{j=\infty} \{j(j+1)c_{j+1} + 2(l+1)(j+1)c_{j+1} - \sqrt{\varepsilon}jc_j + [\lambda - \sqrt{\varepsilon}(l+1)]c_j\}\rho^j = 0,$$

$$(j+1)(2l+j+2)c_{j+1} + [\lambda - \sqrt{\varepsilon}(l+j+1)]c_j = 0$$

$$c_{j+1} = \frac{\sqrt{\varepsilon}(l+j+1) - \lambda}{(j+1)(2l+j+2)}c_j$$

The solution is

$$L(\rho) = c_0 \left\{ 1 + \frac{\sqrt{\varepsilon}(l+1) - \lambda}{(2l+2)}\rho + \frac{(\sqrt{\varepsilon}(l+1) - \lambda)(\sqrt{\varepsilon}(l+2) - \lambda)}{(2l+2)2(2l+3)}\rho^2 + \right. \\ \left. + \frac{(\sqrt{\varepsilon}(l+1) - \lambda)(\sqrt{\varepsilon}(l+2) - \lambda)(\sqrt{\varepsilon}(l+3) - \lambda)}{(2l+2)2(2l+3)3(2l+4)}\rho^3 + \dots \right\}$$

To keep the solution from diverging at $\rho \rightarrow \infty$, the series terms vanish for

$$\frac{\lambda}{\sqrt{\varepsilon}} = l+1, l+2, l+3, \dots,$$

and we obtain the Laguerre Polynomials.

A solution for $R(\rho)$ is the infinite linear combination

$$\alpha_0 L_0(\rho)\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + \alpha_1 L_1(\rho)\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + \alpha_2 L_2(\rho)\rho^l e^{-\frac{1}{2}\sqrt{\varepsilon}\rho} + \dots$$

0.4 The Laguerre Series Associated with $f(x)$

Let $f(x)$ be defined on $[0, \infty)$, and let $L_n(x)$ be the Laguerre

Polynomials

$$L_0(x) = 1, L_1(x) = -x + 1, L_2(x) = \frac{1}{2}x^2 - 2x + 1,$$

$$H_3(x) = -\frac{1}{6}x^3 + \frac{3}{2}x^2 - 3x + 1, \dots$$

The Polynomials are orthogonal on $[0, \infty)$. That is,

$$\int_{x=0}^{x=\infty} e^{-x} L_m(x) L_n(x) dx = \delta_{mn}$$

If $f(x)$ can be expanded in the $L_n(x)$,

$$f(x) = a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots,$$

Then,

$$\begin{aligned} \int_{x=0}^{x=\infty} e^{-x} f(x) L_n(x) dx &= \int_{x=0}^{x=\infty} e^{-x} \{a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots\} L_n(x) dx \\ &= a_0 \underbrace{\int_{x=0}^{x=\infty} e^{-x} L_0(x) L_n(x) dx}_{\delta_{0n}} + a_1 \underbrace{\int_{x=0}^{x=\infty} e^{-x} L_1(x) L_n(x) dx}_{\delta_{1n}} + a_2 \underbrace{\int_{x=0}^{x=\infty} e^{-x} L_2(x) L_n(x) dx}_{\delta_{2n}} + \dots \\ &= a_n. \end{aligned}$$

Thus, the Laguerre coefficients are

$$a_n = \int_{\xi=-\infty}^{\xi=\infty} e^{-\xi} f(\xi) L_n(\xi) d\xi.$$

The Laguerre Series associated with $f(x)$ is

$$a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots$$

1.

Divergence of the Laguerre Kernel in the Calculus of Limits

Calculus of Limits Conditions for the Laguerre Series to equal its function reflect the belief that a smooth enough function equals its Laguerre Series.

In fact, in the Calculus of Limits, no smoothness of the function guarantees even the convergence of the Laguerre Series.

In the Calculus of Limits, the Laguerre Series is the limit of the sequence of Partial Sums

$$\begin{aligned} \mathcal{L}_{\text{Laguerre}} \mathcal{S}_n \{f(x)\} &= a_0 L_0(x) + \dots + a_n L_n(x) \\ &= \left(\int_{\xi=0}^{\xi=\infty} e^{-x} f(\xi) L_0(\xi) d\xi \right) L_0(x) + \dots + \left(\int_{\xi=0}^{\xi=\infty} e^{-x} f(\xi) L_n(\xi) d\xi \right) L_n(x) \\ &= \int_{\xi=0}^{\xi=\infty} f(\xi) e^{-x} \{L_0(\xi) L_0(x) + \dots + L_n(\xi) L_n(x)\} d\xi. \end{aligned}$$

As $n \rightarrow \infty$, the Laguerre Sequence

$$e^{-x} \{L_0(\xi) L_0(x) + \dots + L_n(\xi) L_n(x)\}$$

becomes the Laguerre Kernel,

$$e^{-x} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \},$$

To see that it is singular at $\xi = x$, we apply Christoffel's Summation Formula for Laguerre's Polynomials, [Spanier, p.212],

$$L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) = (n + 1) \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x}.$$

For $\xi \rightarrow x$,

$$e^{-x} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) \} \rightarrow e^{-x} \{ L_0^2(x) + \dots + L_n^2(x) \},$$

and

$$e^{-x}(n + 1) \frac{L_{n+1}(\xi)L_n(x) - L_n(\xi)L_{n+1}(x)}{\xi - x} \rightarrow e^{-x}(n + 1) \frac{0}{0}.$$

Applying Bernoulli's rule to the indeterminate limit,

$$\begin{aligned} \lim_{\xi \rightarrow x} \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x} &= \lim_{\xi \rightarrow x} \frac{D_\xi L_n(\xi)L_{n+1}(x) - D_\xi L_{n+1}(\xi)L_n(x)}{D_\xi(\xi - x)} \\ &= \lim_{\xi \rightarrow x} [L_n'(\xi)L_{n+1}(x) - L_{n+1}'(\xi)L_n(x)] \\ &= L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x) \end{aligned}$$

Therefore,

$$e^{-x} \{ L_0^2(x) + \dots + L_n^2(x) \} = (n + 1)[L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x)].$$

Since $\varphi_n(x)$, and $\varphi_{n+1}(x)$ solve the differential equation, [Szegő, p.99, #5.1.2],

$$\underbrace{x}_{a(x)} \cdot y''(x) + \underbrace{(1-x)}_{b(x)} \cdot y'(x) + ny(x) = 0,$$

we have,

$$\begin{aligned}
L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x) &= (\text{const})e^{-\int \frac{b(x)}{a(x)}dx} \\
&= (\text{const})e^{\int (1-\frac{1}{x})dx} \\
&= (\text{const})e^{x-\log x}.
\end{aligned}$$

Hence,

$$\begin{aligned}
e^{-x} \{L_0^2(x) + \dots + L_n^2(x)\} &= e^{-x}(n+1)(\text{const})e^{x-\log x} \\
&= (n+1)(\text{const})e^{-\log x} \\
&= (n+1)(\text{const})\frac{1}{x}.
\end{aligned}$$

for any $0 < x < \infty$.

Thus, as $n \rightarrow \infty$, Laguerre's Kernel diverges to ∞ at any $\xi = x$.

Therefore, while the partial sums of the Laguerre Series exist, their limit does not. That is, due to the singularity at $\xi = x$, the Laguerre Series does not converge in the Calculus of Limits.

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because at any $\xi \neq x$, the Laguerre Kernel vanishes, and the integral will be identically zero, for any function $f(x)$.

To see that the kernel vanishes for any $\xi \neq x$, we apply the Christoffel Summation Formula with $\xi \neq x$,

$$L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) = (n+1) \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x}.$$

By Rodrigue's Formula [Szego2, p.100]

$$L_n(x) = e^x \frac{1}{n!} D_x^n \{ e^{-x} x^n \}$$

Thus,

$$\begin{aligned} e^{-x} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) \} &= \\ &= e^{-x} (n+1) \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x} \\ &= \frac{e^\xi}{\xi - x} \underbrace{\frac{n+1}{n!(n+1)!}}_{\rightarrow 0, n \rightarrow \infty} \underbrace{\{ D_\xi^n \{ e^{-\xi} \xi^n \} D_x^{n+1} \{ e^{-x} x^n \} - D_\xi^{n+1} \{ e^{-\xi} \xi^n \} D_x^n \{ e^{-x} x^n \} \}}_{\rightarrow 0, n \rightarrow \infty} \\ &\rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

That is, the Laguerre Kernel vanishes for any $\xi \neq x$.


Plots of the Laguerre Sequence confirm that

In the Calculus of Limits,

the Laguerre Kernel is either singular or zero


1.2 Plots of $e^{-x} \{ L_0(\xi)L_0(x) + L_1(\xi)L_1(x) + L_2(\xi)L_2(x) \}$

In Maple, $\text{plot}(\sum_{i=0}^{365} e^{-x} \text{LaguerreL}(i, 7)) * \text{LaguerreL}(i, x), x = .1 .. 17)$



The oscillations away from the origin, are suppressed more effectively. However, a singularity close to the origin, which is the case of an electron orbiting a proton, is effectively described

In Maple, $\text{plot}\left(\sum_{i=0}^{223} e^{-x} \text{LaguerreL}(i, 0.2)\right) * \text{LaguerreL}(i, x), x = .1 .. 23$



The plots confirm that, the Laguerre Series Theorem cannot be proved in the Calculus of Limits.

1.3 Infinitesimal Calculus Solution

By resolving the problem of the infinitesimals [Dan2], we obtained the Infinite Hyper-reals that are strictly smaller than ∞ , and constitute the value of the Delta Function at the singularity.

The controversy surrounding the Leibnitz Infinitesimals derailed the development of the Infinitesimal Calculus, and the Delta Function could not be defined and investigated properly.

In Infinitesimal Calculus, [Dan3], we can differentiate over jump discontinuities, and integrate over singularities.

The Delta Function, the idealization of an impulse in Radar circuits, is a Discontinuous Hyper-Real function which definition requires Infinite Hyper-reals, and which analysis requires Infinitesimal Calculus.

In [Dan5], we show that in infinitesimal Calculus, the hyper-real

$$\delta(x) = \frac{1}{2\pi} \int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega$$

is zero for any $x \neq 0$,

it spikes at $x = 0$, so that its Infinitesimal Calculus

$$\text{integral is } \int_{x=-\infty}^{x=\infty} \delta(x)dx = 1,$$

$$\text{and } \delta(0) = \frac{1}{dx} < \infty.$$

Here, we show that in Infinitesimal calculus, the Laguerre Kernel is a hyper-real Delta Function.

And the Laguerre Series $\mathcal{L}_{egendre} \mathcal{S} \{f(x)\}$ associated with a Hyper-real function $f(x)$, equals $f(x)$.

2.

Hyper-real Line

Each real number α can be represented by a Cauchy sequence of rational numbers, (r_1, r_2, r_3, \dots) so that $r_n \rightarrow \alpha$.

The constant sequence $(\alpha, \alpha, \alpha, \dots)$ is a constant hyper-real.

In [Dan2] we established that,

1. Any totally ordered set of positive, monotonically decreasing to zero sequences (l_1, l_2, l_3, \dots) constitutes a family of infinitesimal hyper-reals.
2. The infinitesimals are smaller than any real number, yet strictly greater than zero.
3. Their reciprocals $\left(\frac{1}{l_1}, \frac{1}{l_2}, \frac{1}{l_3}, \dots\right)$ are the infinite hyper-reals.
4. The infinite hyper-reals are greater than any real number, yet strictly smaller than infinity.
5. The infinite hyper-reals with negative signs are smaller than any real number, yet strictly greater than $-\infty$.
6. The sum of a real number with an infinitesimal is a non-constant hyper-real.
7. The Hyper-reals are the totality of constant hyper-reals, a family of infinitesimals, a family of infinitesimals with

- negative sign, a family of infinite hyper-reals, a family of infinite hyper-reals with negative sign, and non-constant hyper-reals.
8. The hyper-reals are totally ordered, and aligned along a line: the Hyper-real Line.
 9. That line includes the real numbers separated by the non-constant hyper-reals. Each real number is the center of an interval of hyper-reals, that includes no other real number.
 10. In particular, zero is separated from any positive real by the infinitesimals, and from any negative real by the infinitesimals with negative signs, $-dx$.
 11. Zero is not an infinitesimal, because zero is not strictly greater than zero.
 12. We do not add infinity to the hyper-real line.
 13. The infinitesimals, the infinitesimals with negative signs, the infinite hyper-reals, and the infinite hyper-reals with negative signs are semi-groups with respect to addition. Neither set includes zero.
 14. The hyper-real line is embedded in \mathbb{R}^∞ , and is not homeomorphic to the real line. There is no bi-continuous one-one mapping from the hyper-real onto the real line.

15. In particular, there are no points on the real line that can be assigned uniquely to the infinitesimal hyper-reals, or to the infinite hyper-reals, or to the non-constant hyper-reals.
16. No neighbourhood of a hyper-real is homeomorphic to an \mathbb{R}^n ball. Therefore, the hyper-real line is not a manifold.
17. The hyper-real line is totally ordered like a line, but it is not spanned by one element, and it is not one-dimensional.

3.

Integral of a Hyper-real Function

In [Dan3], we defined the integral of a Hyper-real Function.

Let $f(x)$ be a hyper-real function on the interval $[a, b]$.

The interval may not be bounded.

$f(x)$ may take infinite hyper-real values, and need not be bounded.

At each

$$a \leq x \leq b,$$

there is a rectangle with base $[x - \frac{dx}{2}, x + \frac{dx}{2}]$, height $f(x)$, and area

$$f(x)dx.$$

We form the **Integration Sum** of all the areas for the x 's that start at $x = a$, and end at $x = b$,

$$\sum_{x \in [a, b]} f(x)dx.$$

If for any infinitesimal dx , the Integration Sum has the same hyper-real value, then $f(x)$ is integrable over the interval $[a, b]$.

Then, we call the Integration Sum the integral of $f(x)$ from $x = a$, to $x = b$, and denote it by

$$\int_{x=a}^{x=b} f(x)dx .$$

If the hyper-real is infinite, then it is the integral over $[a, b]$,

If the hyper-real is finite,

$$\int_{x=a}^{x=b} f(x)dx = \text{real part of the hyper-real.} \square$$

3.1 The countability of the Integration Sum

In [Dan1], we established the equality of all positive infinities:

We proved that the number of the Natural Numbers,

$Card\mathbb{N}$, equals the number of Real Numbers, $Card\mathbb{R} = 2^{Card\mathbb{N}}$, and

we have

$$Card\mathbb{N} = (Card\mathbb{N})^2 = \dots = 2^{Card\mathbb{N}} = 2^{2^{Card\mathbb{N}}} = \dots \equiv \infty .$$

In particular, we demonstrated that the real numbers may be well-ordered.

Consequently, there are countably many real numbers in the interval $[a, b]$, and the Integration Sum has countably many terms.

While we do not sequence the real numbers in the interval, the summation takes place over countably many $f(x)dx$.

The Lower Integral is the Integration Sum where $f(x)$ is replaced

by its lowest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.2} \quad \sum_{x \in [a, b]} \left(\inf_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

The Upper Integral is the Integration Sum where $f(x)$ is replaced

by its largest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.3} \quad \sum_{x \in [a, b]} \left(\sup_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

If the integral is a finite hyper-real, we have

3.4 *A hyper-real function has a finite integral if and only if its upper integral and its lower integral are finite, and differ by an infinitesimal.*

4.

Delta Function

In [Dan5], we have defined the Delta Function, and established its properties

1. The Delta Function is a hyper-real function defined from the

hyper-real line into the set of two hyper-reals $\left\{0, \frac{1}{dx}\right\}$. The

hyper-real 0 is the sequence $\langle 0, 0, 0, \dots \rangle$. The infinite hyper-

real $\frac{1}{dx}$ depends on our choice of dx .

2. We will usually choose the family of infinitesimals that is

spanned by the sequences $\left\langle \frac{1}{n} \right\rangle, \left\langle \frac{1}{n^2} \right\rangle, \left\langle \frac{1}{n^3} \right\rangle, \dots$. It is a

semigroup with respect to vector addition, and includes all

the scalar multiples of the generating sequences that are

non-zero. That is, the family includes infinitesimals with

negative sign. Therefore, $\frac{1}{dx}$ will mean the sequence $\langle n \rangle$.

Alternatively, we may choose the family spanned by the

sequences $\left\langle \frac{1}{2^n} \right\rangle, \left\langle \frac{1}{3^n} \right\rangle, \left\langle \frac{1}{4^n} \right\rangle, \dots$. Then, $\frac{1}{dx}$ will mean the

sequence $\langle 2^n \rangle$. Once we determined the basic infinitesimal dx , we will use it in the Infinite Riemann Sum that defines an Integral in Infinitesimal Calculus.

3. The Delta Function is strictly smaller than ∞

4. We define, $\delta(x) \equiv \frac{1}{dx} \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x)$,

$$\text{where } \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x) = \begin{cases} 1, & x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right] \\ 0, & \text{otherwise} \end{cases}.$$

5. Hence,

$$\diamond \text{ for } x < 0, \delta(x) = 0$$

$$\diamond \text{ at } x = -\frac{dx}{2}, \delta(x) \text{ jumps from } 0 \text{ to } \frac{1}{dx},$$

$$\diamond \text{ for } x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right], \delta(x) = \frac{1}{dx}.$$

$$\diamond \text{ at } x = 0, \delta(0) = \frac{1}{dx}$$

$$\diamond \text{ at } x = \frac{dx}{2}, \delta(x) \text{ drops from } \frac{1}{dx} \text{ to } 0.$$

$$\diamond \text{ for } x > 0, \delta(x) = 0.$$

$$\diamond x\delta(x) = 0$$

6. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \left\langle \mathcal{X}_{[-\frac{1}{2}, \frac{1}{2}]}(x), 2\mathcal{X}_{[-\frac{1}{4}, \frac{1}{4}]}(x), 3\mathcal{X}_{[-\frac{1}{6}, \frac{1}{6}]}(x), \dots \right\rangle$

7. If $dx = \langle \frac{2}{n} \rangle$, $\delta(x) = \left\langle \frac{1}{2 \cosh^2 x}, \frac{2}{2 \cosh^2 2x}, \frac{3}{2 \cosh^2 3x}, \dots \right\rangle$

8. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \langle e^{-x}\chi_{[0,\infty)}, 2e^{-2x}\chi_{[0,\infty)}, 3e^{-3x}\chi_{[0,\infty)}, \dots \rangle$

9.
$$\int_{x=-\infty}^{x=\infty} \delta(x)dx = 1.$$

10.
$$\delta(\xi - x) = \frac{1}{2\pi} \int_{k=-\infty}^{k=\infty} e^{-ik(\xi-x)}dk$$

5.

Convergent Series

In [Dan8], we defined convergence of infinite series in Infinitesimal Calculus

5.1 Sequence Convergence to a finite hyper-real a

$$a_n \rightarrow a \text{ iff } \langle a_n \rangle - \langle a \rangle = \text{infinitesimal.}$$

5.2 Sequence Convergence to an infinite hyper-real A

$$a_n \rightarrow A \text{ iff } \langle a_n \rangle \text{ represents the infinite hyper-real } A.$$

5.3 Series Convergence to a finite hyper-real s

$$a_1 + a_2 + \dots \rightarrow s \text{ iff } \langle a_1 + \dots + a_n \rangle - \langle s \rangle = \text{infinitesimal.}$$

5.4 Series Convergence to an Infinite Hyper-real S

$$a_1 + a_2 + \dots \rightarrow S \text{ iff}$$

$$\langle a_1 + \dots + a_n \rangle \text{ represents the infinite hyper-real } S.$$

6.

Laguerre Sequence and $\delta(\xi - x)$

6.1 Laguerre Sequence Definition

The Laguerre Series partial sums

$$\mathcal{L}_{\text{Laguerre}} \mathcal{S}_n \{f(x)\} = \int_{\xi=0}^{\xi=\infty} f(\xi) e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} d\xi.$$

give rise to the Laguerre Sequence

$$L_n(\xi, x) = e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\}.$$

6.2 Laguerre Sequence is a Delta Sequence

For each $n = 0, 1, 2, 3, \dots$

$$L_n(\xi, x) = e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\},$$

1. *has the sifting property*

$$\int_{\xi=0}^{\xi=\infty} e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} d\xi = 1$$

2. *is a continuous function*

3. *peaks for each $\xi \rightarrow x$ to $\lim_{\xi \rightarrow x} L_n(\xi, x) \geq (n + 1) \text{const.}$*

Proof of (1)

$$\begin{aligned}
& \int_{\xi=0}^{\xi=\infty} e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} d\xi = \\
& = \underbrace{L_0(x)}_1 \underbrace{\int_{\xi=0}^{\xi=\infty} e^{-\xi} \underbrace{L_0(\xi)}_1 d\xi}_1 + L_1(x) \int_{\xi=0}^{\xi=\infty} e^{-\xi} L_1(\xi) d\xi + \dots + L_n(x) \int_{\xi=0}^{\xi=\infty} e^{-\xi} L_n(\xi) d\xi
\end{aligned}$$

By [Spiegel, p.154, #28.22], for $k = 1, 2, \dots, n$,

$$\int_{\xi=0}^{\xi=\infty} x^p e^{-x} L_k(\xi) d\xi = \begin{cases} 0, & p < k \\ (-1)^n k!, & p = k \end{cases}$$

Therefore, for $k = 1, 2, \dots, n$,

$$\int_{\xi=0}^{\xi=\infty} e^{-x} L_k(\xi) d\xi = 0.$$

Hence,

$$\int_{\xi=0}^{\xi=\infty} e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} d\xi = 1,$$

That is, the Laguerre Sequence has the sifting property. \square

Proof of (3)

By Christoffel's Summation Formula for Laguerre's Polynomials,

[Spanier, p.212],

$$L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) = (n+1) \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x}.$$

For $\xi \rightarrow x$,

$$e^{-x} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} \rightarrow e^{-x} \{L_0^2(x) + \dots + L_n^2(x)\},$$

and

$$e^{-x}(n+1) \frac{L_{n+1}(\xi)L_n(x) - L_n(\xi)L_{n+1}(x)}{\xi - x} \rightarrow e^{-x}(n+1) \frac{0}{0}.$$

Applying Bernoulli's rule to the indeterminate limit,

$$\begin{aligned} \lim_{\xi \rightarrow x} \frac{L_n(\xi)L_{n+1}(x) - L_{n+1}(\xi)L_n(x)}{\xi - x} &= \lim_{\xi \rightarrow x} \frac{D_\xi L_n(\xi)L_{n+1}(x) - D_\xi L_{n+1}(\xi)L_n(x)}{D_\xi(\xi - x)} \\ &= \lim_{\xi \rightarrow x} [L_n'(\xi)L_{n+1}(x) - L_{n+1}'(\xi)L_n(x)] \\ &= L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x) \end{aligned}$$

Therefore,

$$e^{-x} \{L_0^2(x) + \dots + L_n^2(x)\} = (n+1)[L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x)].$$

Since $\varphi_n(x)$, and $\varphi_{n+1}(x)$ solve the differential equation, [Szegő2, p.99, #5.1.2],

$$\underbrace{x}_{a(x)} \cdot y''(x) + \underbrace{(1-x)}_{b(x)} \cdot y'(x) + ny(x) = 0,$$

we have,

$$\begin{aligned} L_n'(x)L_{n+1}(x) - L_{n+1}'(x)L_n(x) &= (const)e^{-\int \frac{b(x)}{a(x)} dx} \\ &= (const)e^{\int (1-\frac{1}{x}) dx} \\ &= (const)e^{x-\log x}. \end{aligned}$$

Hence,

$$e^{-x} \{L_0^2(x) + \dots + L_n^2(x)\} = e^{-x}(n+1)(const)e^{x-\log x}$$

$$= (n + 1)(const)e^{-\log x}$$

$$= (n + 1)(const)\frac{1}{x}$$

for any $0 < x < \infty$. \square

7.

Laguerre Kernel and $\delta(\xi - x)$

7.1 Laguerre Kernel in the Calculus of Limits

The Laguerre Series partial sums

$$\mathcal{L}_{Laguerre} \mathcal{S}_n \{f(x)\} = \int_{\xi=0}^{\xi=\infty} f(\xi) e^{-\xi} \underbrace{\{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\}}_{\text{Laguerre Sequence}} d\xi.$$

give rise to the Laguerre Sequence.

The limit of the Laguerre Sequence is an infinite series called the Laguerre Kernel

$$\mathcal{L}_{Laguerre}(\xi - x) = e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots\}$$

7.2 *In the Calculus of Limits, the Laguerre Kernel does not have the sifting property*

Proof: for $\xi \rightarrow x$,

$$e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x)\} \geq \text{const} \cdot (n+1) \frac{1}{x}$$

$$\xrightarrow[n \rightarrow \infty]{} \infty$$

That is, for $\xi \rightarrow x$,

$e^{-\xi} \{L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots\}$ is singular. \square

7.3 Hyper-real Laguerre Kernel in Infinitesimal Calculus

$$\begin{aligned}
 \mathcal{L}_{\text{Laguerre}}(\xi - x) &= e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \} \\
 &= \begin{cases} \langle n \rangle, & \xi = x \\ 0, & \xi \neq x \end{cases} \\
 &= \delta(\xi - x).
 \end{aligned}$$

Proof: $\mathcal{L}_{\text{Laguerre}}(\xi - x) = e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \}$

$$= \begin{cases} \langle n \rangle, & \xi = x \\ 0, & \xi \neq x \end{cases}$$

Denoting by $\frac{1}{dx}$ the infinite hyper-real $\langle n \rangle$,

$$\begin{aligned}
 &= \begin{cases} 0, & \xi \neq x \\ \frac{1}{dx}, & \xi = x \end{cases} \\
 &= \delta(\xi - x). \square
 \end{aligned}$$

8.

Laguerre Series and $\delta(\xi - x)$

8.1 Laguerre Series of a Hyper-real Function

Let $f(x)$ be a hyper-real function integrable on $[0, \infty)$.

Then, for each $n = 0, 1, 2, 3, \dots$, the integrals

$$a_n = \int_{x=0}^{x=\infty} e^{-x} f(x) L_n(x) dx$$

exist, with finite, or infinite hyper-real values. The a_n are the Laguerre Coefficients of $f(x)$.

The Laguerre Series associated with $f(x)$ is

$$\mathcal{L}_{\text{Laguerre}} \mathcal{S} \{ f(x) \} = a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots$$

For each x , it may assume finite or infinite hyper-real values.

8.2

$$\mathcal{L}_{\text{Laguerre}} \mathcal{S} \{ \delta(\xi - x) \} = \delta(\xi - x)$$

Proof:

$$\mathcal{L}_{\text{Laguerre}} \mathcal{S} \{ \delta(\xi - x) \} = a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots$$

where

$$a_n = \int_{x=0}^{x=\infty} e^{-x} \delta(\xi - x) L_n(x) dx.$$

$$= e^{-\xi} L_n(\xi)$$

Therefore,

$$\begin{aligned} \mathcal{L}_{\text{aguerre}} \mathcal{S} \{ \delta(\xi - x) \} &= e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \} \\ &= \mathcal{L}_{\text{aguerre}}(\xi - x), \text{ by 7.3,} \\ &= \delta(\xi - x), \text{ by 7.3.} \square \end{aligned}$$

9.

Laguerre Series Theorem

The Laguerre Series Theorem for a hyper-real function, $f(x)$, is the Fundamental Theorem of Laguerre Series.

It supplies the conditions under which the Laguerre Series associated with $f(x)$ equals $f(x)$.

It is believed to hold in the Calculus of Limits under the Picone Conditions, or under the Hobson Conditions [Sansone]. In fact,

The Theorem cannot be proved in the Calculus of Limits under any conditions,

because the summation of the Laguerre Series requires integration of the singular Laguerre Kernel.

9.1 Laguerre Series Theorem cannot be proved in the Calculus of Limits

Proof: Let $f(x)$ be integrable on $[0, \infty)$.

In the Calculus of Limits, the Laguerre Series is the limit of the sequence of Partial Sums

$$\mathcal{L}_{\text{Laguerre}} \mathcal{S}_n \{f(x)\} = a_0 L_0(x) + \dots + a_n L_n(x)$$

$$\begin{aligned}
&= \left(\int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi}L_0(\xi)d\xi \right) L_0(x) + \dots + \left(\int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi}L_n(\xi)d\xi \right) L_n(x) \\
&= \int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) \} d\xi .
\end{aligned}$$

As $n \rightarrow \infty$, the Laguerre Sequence

$$e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) \}$$

becomes the Laguerre Kernel,

$$e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \}.$$

By 7.2, the Laguerre Kernel diverges to infinity at any $\xi = x$.

Therefore, while the partial sums of the Laguerre Series exist, their limit does not. Conditions by Uspensky [Sansone] failed to comprehend the sifting through the values of $f(\xi)$ by the Laguerre Kernel, and the picking of $f(\xi)$ at $\xi = x$.

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because for any $\xi \neq x$, the Laguerre Kernel vanishes, and the integral is identically zero, for any function $f(x)$.

Thus, the Laguerre Series Theorem cannot be proved in the Calculus of Limits. \square

9.2 Calculus of Limits Conditions are insufficient and irrelevant for the Laguerre Series Theorem

Proof: The Uspensky Conditions [Sansone, p.384] are

1. $f(x)$ integrable in any $[a, b]$ in $[0, \infty)$, and
of bounded variation in $|\xi - x| < \delta$, for any $0 < \delta < x$
2. $e^{-x} f^2(x)$ integrable on $[\gamma, \infty)$ for any $\gamma > 0$
3. $e^{-x} f^2(x)$ integrable on $[0, \beta]$ for any $\beta > 0$

It is clear from 9.1 that these conditions on $f(x)$, that have nothing to do with the Laguerre Polynomials, do not resolve the singularity of the Laguerre kernel, and are insufficient for, and irrelevant to the Laguerre Series Theorem. \square

In Infinitesimal Calculus, by 7.3, the Laguerre Kernel is the Delta Function, and by 8.2, it equals its Laguerre Series.

Then, the Laguerre Series Theorem holds for any Hyper-Real Function:

9.3 Laguerre Series Theorem for Hyper-real $f(x)$

If $f(x)$ is hyper-real function integrable on $(-\infty, \infty)$

Then,
$$f(x) = \mathcal{L}_{\text{aguerre}} \mathcal{S} \{ f(x) \}$$

Proof:

$$f(x) = \int_{\xi=0}^{\xi=\infty} f(\xi)\delta(\xi - x)d\xi$$

Substituting from 7.3,

$$\delta(\xi - x) = e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \},$$

$$f(x) = \int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \} d\xi$$

This Hyper-real Integral is the summation,

$$\sum_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi} \{ L_0(\xi)L_0(x) + \dots + L_n(\xi)L_n(x) + \dots \} d\xi$$

which amounts to the hyper-real function $f(x)$, and is well-defined.

Hence, the summation of each term in the integrand exists, and we may write the integral as the sum

$$\begin{aligned} &= \underbrace{\left(\int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi^2} L_0(\xi)d\xi \right)}_{a_0} L_0(x) + \dots + \underbrace{\left(\int_{\xi=0}^{\xi=\infty} f(\xi)e^{-\xi} L_n(\xi)d\xi \right)}_{a_n} L_n(x) + \dots \\ &= a_0 L_0(x) + a_1 L_1(x) + a_2 L_2(x) + \dots \\ &= \mathcal{L}_{aguerre} \mathcal{S} \{ f(x) \}. \square \end{aligned}$$

In particular, the Delta Function violates the Calculus of Limits Conditions

❖ *The Hyper-real $\delta(x)$, is not defined in the Calculus of Limits,
and is not integrable in any interval.*

But by 8.2, $\delta(\xi - x)$ satisfies the Laguerre Series Theorem.

References

[Abramowitz] Abramowitz, M., and Stegun, I., “*Handbook of Mathematical Functions with Formulas Graphs and Mathematical Tables*”, U.S. Department of Commerce, National Bureau of Standards, 1964.

[Dan1] Dannon, H. Vic, “*Well-Ordering of the Reals, Equality of all Infinities, and the Continuum Hypothesis*” in Gauge Institute Journal Vol. 6 No. 2, May 2010;

[Dan2] Dannon, H. Vic, “*Infinitesimals*” in Gauge Institute Journal Vol.6 No. 4, November 2010;

[Dan3] Dannon, H. Vic, “*Infinitesimal Calculus*” in Gauge Institute Journal Vol. 7 No. 4, November 2011;

[Dan4] Dannon, H. Vic, “*Riemann’s Zeta Function: the Riemann Hypothesis Origin, the Factorization Error, and the Count of the Primes*”, in Gauge Institute Journal of Math and Physics, Vol. 5, No. 4, November 2009.

[Dan5] Dannon, H. Vic, “*The Delta Function*” in Gauge Institute Journal Vol. 8, No. 1, February, 2012;

[Dan6] Dannon, H. Vic, “*Riemannian Trigonometric Series*”, Gauge Institute Journal, Volume 7, No. 3, August 2011.

[Dan7] Dannon, H. Vic, “*Delta Function the Fourier Transform, and the Fourier Integral Theorem*” in Gauge Institute Journal Vol. 8, No. 2, May, 2012;

[Dan8] Dannon, H. Vic, “*Infinite Series with Infinite Hyper-real Sum* ” in Gauge Institute Journal Vol. 8, No. 3, August, 2012;

[Ferrers] Ferrers, N., M., “*An Elementary treatment on Spherical Harmonics*”, Macmillan, 1877.

[Gradshteyn] Gradshteyn, I., S., and Ryzhik, I., M., “*Tables of Integrals Series and Products*”, 7th Edition, edited by Allan Jeffery, and Daniel Zwillinger, Academic Press, 2007

[Hardy] Hardy, G. H., *Divergent Series*, Chelsea 1991.

[Jackson] Jackson, Dunham, “*Fourier Series and Orthogonal Polynomials*”, Mathematical association of America, 1941.

[Magnus] Magnus, W., Oberhettinger, F., Sony, R., P., “*Formulas and Theorems for the Special Functions of Mathematical Physics*” Third Edition, Springer-Verlag, 1966.

[Sansone] Sansone, Giovanni, “*Orthogonal Functions*”, Revised Edition, Krieger, 1977.

[Spiegel] Spiegel, Murray, “*Mathematical Handbook of formulas and tables*” Schaum’s Outline Series, McGraw Hill, 1968.

[Spanier] Spanier, Jerome, and Oldham, Keith, “*An Atlas of Functions*”, Hemisphere, 1987.

[Szego2] Szego, Gabor, “*Orthogonal Polynomials*” Revised Edition, American Mathematical Society, 1959.

[Szego4] Szego, Gabor, “*Orthogonal Polynomials*” Fourth Edition, American Mathematical Society, 1975.

[Weisstein], Weisstein, Eric, W., “*CRC Encyclopedia of Mathematics*”, Third Edition, CRC Press, 2009.