

Periodic Delta Function, and Poisson Integral for Abel Summation of Fourier Series,

H. Vic Dannon
vic0@comcast.net
June, 2012

Abstract The Poisson Integral Theorem supplies the conditions under which the Poisson Integral at $r = 1 - dr$, associated with a function $f(x)$ equals $f(x)$.

It is believed to hold in the Calculus of Limits under some conditions. In fact,

*The Theorem cannot be proved in the Calculus of Limits
under any conditions,*

because it requires integration of the singular Poisson Kernel.

In Infinitesimal Calculus, the Poisson Kernel

$$\frac{1}{2} \frac{dr(2 - dr)}{(1 - dr)(1 - \cos \pi[\xi - x]) + (dr)^2}$$

is the Periodic Delta Function,

$$\delta_{Periodic}(\xi - x) = \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots$$

$\delta_{Periodic}(\xi - x)$ violates the Calculus of Limits Conditions

- ❖ *The Hyper-real $\delta(x)$, is not defined in the Calculus of Limits, and $|\delta(x)|$ is not integrable in any bounded interval.*
- ❖ *$\frac{1}{2}(\delta(x+0) + \delta(x-0)) = 0$ does not replace $\delta(x)$ at its discontinuity point, $x = 0$.*

But $\delta_{Periodic}(\xi - x)$ equals the Poisson Integral associated with it at $r = 1 - dr$.

The Poisson Integral associated with any hyper-real periodic integrable $f(x)$ at $r = 1 - dr$, equals $f(x)$.

Keywords: Infinitesimal, Infinite-Hyper-Real, Hyper-Real, infinite Hyper-real, Infinitesimal Calculus, Delta Function, Fourier Transform, Periodic Delta Function, Delta Comb, Fourier Series, Poisson Kernel, Poisson Integral

2000 Mathematics Subject Classification 26E35; 26E30; 26E15; 26E20; 26A06; 26A12; 03E10; 03E55; 03E17; 03H15; 46S20; 97I40; 97I30.

Contents

0. The Origin of the Poisson Integral Theorem
1. Divergence of the Poisson Kernel in the Calculus of Limits
2. Hyper-real line.
3. Integral of a Hyper-real Function
4. Delta Function
5. Periodic Delta Function, $\delta_{Periodic}(\xi - x)$
6. Convergent Series
7. Poisson Sequence and $\delta_{Periodic}(\xi - x)$
8. Poisson Kernel and $\delta_{Periodic}(\xi - x)$
9. Poisson Integral and $\delta_{Periodic}(\xi - x)$
10. Poisson Integral Theorem

References

1.

The Origin of Poisson Integral

Theorem

Let $f(x)$ be a function integrable on $[-1,1]$, so that $f(1) = f(-1)$.

The Fourier Coefficients of $f(x)$ are

$$\frac{1}{2} \int_{\xi=-1}^{\xi=1} f(\xi) e^{-in\pi\xi} d\xi \equiv c_n, \quad n = \dots, -2, -1, 0, 1, 2, \dots$$

The Fourier Series associated with $f(x)$ is

$$\begin{aligned} & \dots + c_{-n} e^{i(-n)\pi x} + \dots + c_{-1} e^{i(-1)\pi x} + c_0 + c_1 e^{i(1)\pi x} + \dots + c_n e^{i(n)\pi x} + \dots = \\ & = \int_{\xi=-1}^{\xi=1} f(\xi) \left\{ \dots + \frac{1}{2} e^{-in\pi(\xi-x)} + \dots + \frac{1}{2} e^{-i\pi(\xi-x)} + \frac{1}{2} + \frac{1}{2} e^{i\pi(\xi-x)} + \dots + \frac{1}{2} e^{in\pi(\xi-x)} + \dots \right\} d\xi \end{aligned}$$

0.1 Abel

To assign a numerical value to the divergent series

$$1 - 1 + 1 - 1 + 1 - 1 + \dots,$$

Abel suggested to consider the convergence of

$$1 - r + r^2 - r^3 + r^4 - r^5 + \dots$$

for

$$r = 1 - 0 < 1.$$

For such r , the series is an alternating geometric series with a quotient $q = -r$ that converges to

$$\frac{1}{1-(-r)} = \frac{1}{1+r}.$$

Since

$$\frac{1}{1+r} \Big|_{r=1-0} = \frac{1}{1+1-0} = \frac{1}{2}$$

we conclude that

the infinite series $1 - 1 + 1 - 1 + 1 - 1 + \dots$ has Abel Sum of $\frac{1}{2}$

For any series

$$a_0 + a_1 + a_2 + a_3 + \dots,$$

$$\text{If } a_0 + a_1 r + a_2 r^2 + a_3 r^3 + \dots \Big|_{r=1-0} \rightarrow A,$$

Then $a_0 + a_1 + a_2 + a_3 + \dots$ has Able Sum A

0.2 Poisson

Poisson Applied Abel's summation to Fourier Series.

Then, we consider

$$\begin{aligned} & \lim_{r \uparrow 1} \left\{ \dots + r^n c_{-n} e^{i(-n)\pi x} + \dots + r c_{-1} e^{i(-1)\pi x} + c_0 + r c_1 e^{i(1)\pi x} + \dots + r^n c_n e^{i(n)\pi x} + \dots \right\} = \\ & = \lim_{r \uparrow 1} \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \left\{ \dots + r^n e^{-in\pi(\xi-x)} + \dots + r e^{-i\pi(\xi-x)} + 1 + r e^{i\pi(\xi-x)} + \dots + r^n e^{in\pi(\xi-x)} + \dots \right\} d\xi \\ & = \int_{\xi=-1}^{\xi=1} f(\xi) \lim_{r \uparrow 1} \frac{1}{2} \left\{ \dots + r^n e^{-in\pi(\xi-x)} + \dots + r e^{-i\pi(\xi-x)} + 1 + r e^{i\pi(\xi-x)} + \dots + r^n e^{in\pi(\xi-x)} + \dots \right\} d\xi \end{aligned}$$

To evaluate the Kernel in $\lim_{r \uparrow 1} \{ \}$, note that for

$$z = r e^{i\varphi}, \text{ with } r < 1,$$

$$\frac{1}{2} + r e^{i\varphi} + r^2 e^{2i\varphi} + \dots = \frac{1}{2} + z + z^2 + \dots$$

$$\begin{aligned}
&= \frac{1}{2} + z(1 + z + z^2 + \dots) \\
&= 1 + z \frac{1}{1-z} \\
&= \frac{1}{2} \frac{1+z}{1-z} \\
&= \frac{1}{2} \frac{1 + re^{i\varphi}}{1 - re^{i\varphi}} \\
&= \frac{1}{2} \frac{1 + re^{i\varphi}}{1 - re^{i\varphi}} \frac{1 - re^{-i\varphi}}{1 - re^{-i\varphi}} \\
&= \frac{1}{2} \frac{1 - r^2 + 2ir \sin \varphi}{1 + r^2 - 2r \cos \varphi} \\
&= \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \varphi} + i \frac{r \sin \varphi}{1 + r^2 - 2r \cos \varphi}
\end{aligned}$$

The real part is

$$\frac{1}{2} + r \cos \varphi + r^2 \cos 2\varphi + \dots = \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \varphi}$$

That is,

$$\frac{1}{2} \left\{ \dots + r^2 e^{-2i\varphi} + re^{-i\varphi} + 1 + re^{i\varphi} + r^2 e^{2i\varphi} + \dots \right\} = \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \varphi}$$

Thus, the Fourier Series may have Abel Sum

$$\int_{\xi=-1}^{\xi=1} f(\xi) \lim_{r \uparrow 1} \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \pi(x - \xi)} d\xi$$

The Poisson Integral associated with $f(x)$ at $r = 1 - 0 +$.

The Poisson Integral associated with the function $f(x)$ is clearly different from the Fourier Series associated with $f(x)$, but it may nevertheless converge to $f(x)$.

The equality of

$$\int_{\xi=-1}^{\xi=1} f(\xi) \lim_{r \uparrow 1} \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)} d\xi$$

to $f(x)$ is the Poisson Integral Theorem, and the question is under which conditions does the Theorem hold.

1.

Divergence of the Poisson Kernel in the Calculus of Limits

The limit of the integral is believed to equal $f(x)$ provided that

1. $|f(x)|$ is integrable on $[-1,1]$
2. $f(x)$ is periodic with period $T = 2$
3. $\frac{1}{2}(f(x+0) + f(x-0))$ replaces $f(x)$ at a discontinuity point.

These are the Fejer Conditions stated in [Dan10], for the Fejer Summation Theorem.

These Conditions reflect the belief that the equality depends only on the function, regardless of the singularity of the Poisson Integral Kernel.

In the Calculus of Limits, no smoothness of the function guarantees the convergence of the Poisson Integral.

1.1 The Poisson Integral Kernel is either singular or zero

In the Calculus of Limits,

$$\lim_{r \uparrow 1} \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)} d\xi = \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \lim_{r \uparrow 1} \underbrace{\frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)}}_{\text{Poisson Kernel}} d\xi.$$

As $r \uparrow 1$, the Poisson Sequence becomes the Poisson Kernel, which is singular, and diverges at any $\xi - x = 2m$.

$$x - \xi = 2m \Rightarrow \cos \pi(x - \xi) = 1,$$

$$\begin{aligned} \Rightarrow \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)} &= \frac{1}{2} \frac{1+r}{1-r}, \\ &\xrightarrow{r \uparrow 1} \frac{1}{2} \frac{1+1}{0+} = \infty \end{aligned}$$

Thus, the Poisson Integral diverges in the Calculus of Limits.

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because at any $\xi - x \neq 2m$,

$$\begin{aligned} \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)} &= \frac{1}{2} \frac{1-r^2}{(1-r)^2 + 4r \sin^2[\frac{1}{2}\pi(x-\xi)]} \\ &\xrightarrow{r \uparrow 1} \frac{0}{2 \sin^2[\frac{1}{2}\pi(x-\xi)]} = 0 \end{aligned}$$

That is, the Poisson Kernel vanishes, and the integral is identically zero, for any function $f(x)$.

Plots of the Poisson Sequence confirm that

In the Calculus of Limits,


the Poisson Integral Kernel is either singular or zero

1.2 Plots of Poisson Sequence with $r_k = 1 - \frac{1}{k}$

For $k = 37$,

$$\text{plot}\left(\frac{36}{74 \cdot 36 + 1 - 74 \cdot 36 \cos(\pi x)}, x = -5..5\right)$$


plots the spikes at $x = 0, x = -2, x = 2, x = -4, x = 4$



For $k = 49$,

$$\text{plot}\left(\frac{49}{4901 - 4900 \cdot \cos(\pi x)}, x = -7..7\right)$$

gives 7 spikes



Thus, the Poisson Integral Theorem does not hold in the Calculus of Limits.

1.3 Infinitesimal Calculus Solution

By resolving the problem of the infinitesimals [Dan2], we obtained the Infinite Hyper-reals that are strictly smaller than ∞ , and constitute the value of the Delta Function at the singularity.

The controversy surrounding the Leibnitz Infinitesimals derailed the development of the Infinitesimal Calculus, and the Delta Function could not be defined and investigated properly.

In Infinitesimal Calculus, [Dan3], we can differentiate over jump discontinuities, and integrate over singularities.

The Delta Function, the idealization of an impulse in Radar circuits, is a Discontinuous Hyper-Real function which definition requires Infinite Hyper-reals, and which analysis requires Infinitesimal Calculus.

In [Dan5], we show that in infinitesimal Calculus, the hyper-real

$$\delta(x) = \frac{1}{2\pi} \int_{\omega=-\infty}^{\omega=\infty} e^{i\omega x} d\omega$$

is zero for any $x \neq 0$,

it spikes at $x = 0$, so that its Infinitesimal Calculus

integral is
$$\int_{x=-\infty}^{x=\infty} \delta(x) dx = 1,$$

and
$$\delta(0) = \frac{1}{dx} < \infty.$$

Here, we show that in Infinitesimal calculus, the Poisson Kernel is a periodic hyper-real Delta Function: A periodic train of Delta Functions. And the Poisson Integral associated with a Hyper-real periodic function $f(x)$, at $r = 1 - dr$, equals $f(x)$.

2.

Hyper-real Line

Each real number α can be represented by a Cauchy sequence of rational numbers, (r_1, r_2, r_3, \dots) so that $r_n \rightarrow \alpha$.

The constant sequence $(\alpha, \alpha, \alpha, \dots)$ is a constant hyper-real.

In [Dan2] we established that,

1. Any totally ordered set of positive, monotonically decreasing to zero sequences (l_1, l_2, l_3, \dots) constitutes a family of infinitesimal hyper-reals.
2. The infinitesimals are smaller than any real number, yet strictly greater than zero.
3. Their reciprocals $\left(\frac{1}{l_1}, \frac{1}{l_2}, \frac{1}{l_3}, \dots\right)$ are the infinite hyper-reals.
4. The infinite hyper-reals are greater than any real number, yet strictly smaller than infinity.
5. The infinite hyper-reals with negative signs are smaller than any real number, yet strictly greater than $-\infty$.
6. The sum of a real number with an infinitesimal is a non-constant hyper-real.
7. The Hyper-reals are the totality of constant hyper-reals, a family of infinitesimals, a family of infinitesimals with

- negative sign, a family of infinite hyper-reals, a family of infinite hyper-reals with negative sign, and non-constant hyper-reals.
8. The hyper-reals are totally ordered, and aligned along a line: the Hyper-real Line.
 9. That line includes the real numbers separated by the non-constant hyper-reals. Each real number is the center of an interval of hyper-reals, that includes no other real number.
 10. In particular, zero is separated from any positive real by the infinitesimals, and from any negative real by the infinitesimals with negative signs, $-dx$.
 11. Zero is not an infinitesimal, because zero is not strictly greater than zero.
 12. We do not add infinity to the hyper-real line.
 13. The infinitesimals, the infinitesimals with negative signs, the infinite hyper-reals, and the infinite hyper-reals with negative signs are semi-groups with respect to addition. Neither set includes zero.
 14. The hyper-real line is embedded in \mathbb{R}^∞ , and is not homeomorphic to the real line. There is no bi-continuous one-one mapping from the hyper-real onto the real line.

15. In particular, there are no points on the real line that can be assigned uniquely to the infinitesimal hyper-reals, or to the infinite hyper-reals, or to the non-constant hyper-reals.
16. No neighbourhood of a hyper-real is homeomorphic to an \mathbb{R}^n ball. Therefore, the hyper-real line is not a manifold.
17. The hyper-real line is totally ordered like a line, but it is not spanned by one element, and it is not one-dimensional.

3.

Integral of a Hyper-real Function

In [Dan3], we defined the integral of a Hyper-real Function.

Let $f(x)$ be a hyper-real function on the interval $[a, b]$.

The interval may not be bounded.

$f(x)$ may take infinite hyper-real values, and need not be bounded.

At each

$$a \leq x \leq b,$$

there is a rectangle with base $[x - \frac{dx}{2}, x + \frac{dx}{2}]$, height $f(x)$, and area

$$f(x)dx.$$

We form the **Integration Sum** of all the areas for the x 's that start at $x = a$, and end at $x = b$,

$$\sum_{x \in [a, b]} f(x)dx.$$

If for any infinitesimal dx , the Integration Sum has the same hyper-real value, then $f(x)$ is integrable over the interval $[a, b]$.

Then, we call the Integration Sum the integral of $f(x)$ from $x = a$, to $x = b$, and denote it by

$$\int_{x=a}^{x=b} f(x)dx .$$

If the hyper-real is infinite, then it is the integral over $[a, b]$,

If the hyper-real is finite,

$$\int_{x=a}^{x=b} f(x)dx = \text{real part of the hyper-real. } \square$$

3.1 The countability of the Integration Sum

In [Dan1], we established the equality of all positive infinities:

We proved that the number of the Natural Numbers,

$Card\mathbb{N}$, equals the number of Real Numbers, $Card\mathbb{R} = 2^{Card\mathbb{N}}$, and

we have

$$Card\mathbb{N} = (Card\mathbb{N})^2 = \dots = 2^{Card\mathbb{N}} = 2^{2^{Card\mathbb{N}}} = \dots \equiv \infty .$$

In particular, we demonstrated that the real numbers may be well-ordered.

Consequently, there are countably many real numbers in the interval $[a, b]$, and the Integration Sum has countably many terms.

While we do not sequence the real numbers in the interval, the summation takes place over countably many $f(x)dx$.

The Lower Integral is the Integration Sum where $f(x)$ is replaced

by its lowest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.2} \quad \sum_{x \in [a, b]} \left(\inf_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

The Upper Integral is the Integration Sum where $f(x)$ is replaced

by its largest value on each interval $[x - \frac{dx}{2}, x + \frac{dx}{2}]$

$$\mathbf{3.3} \quad \sum_{x \in [a, b]} \left(\sup_{x - \frac{dx}{2} \leq t \leq x + \frac{dx}{2}} f(t) \right) dx$$

If the integral is a finite hyper-real, we have

3.4 *A hyper-real function has a finite integral if and only if its upper integral and its lower integral are finite, and differ by an infinitesimal.*

4.

Delta Function

In [Dan5], we have defined the Delta Function, and established its properties

1. The Delta Function is a hyper-real function defined from the

hyper-real line into the set of two hyper-reals $\left\{0, \frac{1}{dx}\right\}$. The

hyper-real 0 is the sequence $\langle 0, 0, 0, \dots \rangle$. The infinite hyper-

real $\frac{1}{dx}$ depends on our choice of dx .

2. We will usually choose the family of infinitesimals that is

spanned by the sequences $\left\langle \frac{1}{n} \right\rangle, \left\langle \frac{1}{n^2} \right\rangle, \left\langle \frac{1}{n^3} \right\rangle, \dots$. It is a

semigroup with respect to vector addition, and includes all

the scalar multiples of the generating sequences that are

non-zero. That is, the family includes infinitesimals with

negative sign. Therefore, $\frac{1}{dx}$ will mean the sequence $\langle n \rangle$.

Alternatively, we may choose the family spanned by the

sequences $\left\langle \frac{1}{2^n} \right\rangle, \left\langle \frac{1}{3^n} \right\rangle, \left\langle \frac{1}{4^n} \right\rangle, \dots$. Then, $\frac{1}{dx}$ will mean the

sequence $\langle 2^n \rangle$. Once we determined the basic infinitesimal dx , we will use it in the Infinite Riemann Sum that defines an Integral in Infinitesimal Calculus.

3. The Delta Function is strictly smaller than ∞

4. We define, $\delta(x) \equiv \frac{1}{dx} \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x)$,

$$\text{where } \mathcal{X}_{[-\frac{dx}{2}, \frac{dx}{2}]}(x) = \begin{cases} 1, & x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right] \\ 0, & \text{otherwise} \end{cases}.$$

5. Hence,

$$\diamond \text{ for } x < 0, \delta(x) = 0$$

$$\diamond \text{ at } x = -\frac{dx}{2}, \delta(x) \text{ jumps from } 0 \text{ to } \frac{1}{dx},$$

$$\diamond \text{ for } x \in \left[-\frac{dx}{2}, \frac{dx}{2}\right], \delta(x) = \frac{1}{dx}.$$

$$\diamond \text{ at } x = 0, \delta(0) = \frac{1}{dx}$$

$$\diamond \text{ at } x = \frac{dx}{2}, \delta(x) \text{ drops from } \frac{1}{dx} \text{ to } 0.$$

$$\diamond \text{ for } x > 0, \delta(x) = 0.$$

$$\diamond x\delta(x) = 0$$

6. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \left\langle \mathcal{X}_{[-\frac{1}{2}, \frac{1}{2}]}(x), 2\mathcal{X}_{[-\frac{1}{4}, \frac{1}{4}]}(x), 3\mathcal{X}_{[-\frac{1}{6}, \frac{1}{6}]}(x), \dots \right\rangle$

7. If $dx = \langle \frac{2}{n} \rangle$, $\delta(x) = \left\langle \frac{1}{2 \cosh^2 x}, \frac{2}{2 \cosh^2 2x}, \frac{3}{2 \cosh^2 3x}, \dots \right\rangle$

8. If $dx = \langle \frac{1}{n} \rangle$, $\delta(x) = \langle e^{-x} \chi_{[0, \infty)}, 2e^{-2x} \chi_{[0, \infty)}, 3e^{-3x} \chi_{[0, \infty)}, \dots \rangle$

9.
$$\int_{x=-\infty}^{x=\infty} \delta(x) dx = 1.$$

In [Dan6], we obtained

10.
$$\delta(\xi - x) = \frac{1}{2\pi} \int_{k=-\infty}^{k=\infty} e^{-ik(\xi-x)} dk$$

In [Dan8], we defined the Periodic Delta Function, and obtained

11.

$$\begin{aligned} \delta_{Periodic}(x) &= \dots + \delta(x + 4) + \delta(x + 2) + \delta(x) + \delta(x - 2) + \delta(x - 4) + \dots \\ &= \dots + \frac{1}{2} e^{-in\pi x} + \dots + \frac{1}{2} e^{-i\pi x} + \frac{1}{2} + \frac{1}{2} e^{i\pi x} + \dots + \frac{1}{2} e^{in\pi x} + \dots \end{aligned}$$

5.

Periodic Delta Function $\delta_{Periodic}(\xi - x)$

5.1 Periodic Delta Function

$$\delta_{Periodic}(\xi - x) = \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots$$

is a periodic hyper-real Delta function, with period $T = 2$.

In [Dan8], we obtained

$$\begin{aligned} \delta_{Periodic}(\xi - x) = \dots + \frac{1}{2}e^{-in\pi(\xi-x)} + \dots + \frac{1}{2}e^{-i\pi(\xi-x)} \\ + \frac{1}{2} + \frac{1}{2}e^{i\pi(\xi-x)} + \dots + \frac{1}{2}e^{in\pi(\xi-x)} + \dots \end{aligned}$$

6.

Convergent Series

In [Dan11], we defined convergence of infinite series in Infinitesimal Calculus

6.1 Sequence Convergence to a finite hyper-real a

$$a_n \rightarrow a \text{ iff } \langle a_n \rangle - \langle a \rangle = \text{infinitesimal.}$$

6.2 Sequence Convergence to an infinite hyper-real A

$$a_n \rightarrow A \text{ iff } \langle a_n \rangle \text{ represents the infinite hyper-real } A.$$

6.3 Series Convergence to a finite hyper-real s

$$a_1 + a_2 + \dots \rightarrow s \text{ iff } \langle a_1 + \dots + a_n \rangle - \langle s \rangle = \text{infinitesimal.}$$

6.4 Series Convergence to an Infinite Hyper-real S

$$a_1 + a_2 + \dots \rightarrow S \text{ iff}$$

$$\langle a_1 + \dots + a_n \rangle \text{ represents the infinite hyper-real } S.$$

7.

Poisson Sequence and $\delta_{Periodic}(\xi - x)$

7.1 Poisson Sequence Definition

Let

$$r_k = 1 - \frac{1}{k}, \quad k = 1, 2, 3, \dots$$

The Sequence of Poisson Integrals at $r_k = 1 - \frac{1}{k}$,

$$\begin{aligned} \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(x-\xi)} d\xi \Big|_{r=1-\frac{1}{k}} &= \\ &= \int_{\xi=-1}^{\xi=1} f(\xi) \underbrace{\frac{k-1}{2k(k-1)+1-2k(k-1)\cos \pi(\xi-x)}}_{\text{Poisson Sequence}} d\xi. \end{aligned}$$

gives rise to the Poisson Sequence

$$P_k(\xi - x) = \frac{k-1}{2k(k-1)+1-2k(k-1)\cos \pi(\xi-x)}.$$

7.2 Poisson Sequence is a Periodic Delta Sequence and represents the Periodic Delta Function,

$$\delta_{Periodic}(\xi - x) = \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots$$

$$\text{Each } P_k(x) = \frac{k-1}{2k(k-1)+1-2k(k-1)\cos \pi x}, \quad k=1,2,3,\dots$$

1. *has the sifting property on each interval,*

$$\dots \int_{x=-5}^{x=-3} P_k(x)dx = 1; \quad \int_{x=-3}^{x=-1} P_k(x)dx = 1; \quad \int_{x=-1}^{x=1} P_k(x)dx = 1 \dots$$

2. *is a continuous function*

3. *peaks on each of these interval to* $\lim_{\xi-x \rightarrow 2m} P_k(\xi-x) = k-1$.

Proof of (1)

$$\int_{x=-1}^{x=1} P_k(x)dx = (k-1) \int_{x=-1}^{x=1} \frac{1}{\underbrace{2k(k-1)+1}_p - \underbrace{2k(k-1)\cos \frac{\pi}{a} x}_{-q}} dx$$

By [Spiegel, p.78],

$$\begin{aligned} &= (k-1) \frac{2}{a\sqrt{(p-q)(p+q)}} \arctan \left[\sqrt{\frac{p-q}{p+q}} \tan \frac{1}{2} ax \right]_{x=-1}^{x=1} \\ &= (k-1) \frac{2}{\underbrace{\pi\sqrt{4k(k-1)+1}}_{1/\pi}} \arctan \left[\underbrace{\sqrt{4k(k-1)+1}}_{2k-1} \tan \frac{1}{2} \pi x \right]_{x=-1}^{x=1} \\ &= \frac{1}{\pi} \left\{ \underbrace{\arctan[(2k-1)\tan \frac{1}{2} \pi]}_{\frac{1}{2}\pi} - \underbrace{\arctan[(2k-1)\tan(-\frac{1}{2} \pi)]}_{-\frac{1}{2}\pi} \right\} \\ &= 1. \square \end{aligned}$$

Proof of (3)

As $\xi - x \rightarrow 2m$,

$$\frac{k-1}{2k(k-1)+1-2k(k-1)\cos\pi x} \rightarrow \frac{k-1}{2k(k-1)+1-2k(k-1)} = k-1. \square$$

7.3 Poisson Sequence Represents $\delta_{Periodic}(\xi - x)$

$$\delta_{Periodic}(\xi - x) = \left\langle \frac{k-1}{2k(k-1)[1-\cos\pi(\xi-x)]+1} \right\rangle$$

8.

Poisson Kernel and $\delta_{Periodic}(\xi - x)$

8.1 Poisson Kernel in the Calculus of Limits

The Sequence of Poisson Integrals at $r_k = 1 - \frac{1}{k}$

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1 - r_k^2}{1 + r_k^2 - 2r_k \cos \pi(x - \xi)} d\xi \Bigg|_{r_k=1-\frac{1}{k}} &= \\ &= \int_{\xi=-1}^{\xi=1} f(\xi) \underbrace{\frac{k-1}{2k(k-1) + 1 - 2k(k-1) \cos \pi(x - \xi)}}_{\text{Poisson Sequence}} d\xi. \end{aligned}$$

gives rise to a Poisson Sequence.

Alternatively, we could use $r_k = 1 - \frac{1}{k^2}$, $r_k = 1 - \frac{1}{k^3}, \dots$

Note that the hyper-real $r = 1 - dr$ includes all such sequences.

The limit of the Poisson Sequence is the Poisson Kernel

$$\lim_{k \rightarrow \infty} \frac{k-1}{2k(k-1)[1 - \cos \pi(x - \xi)] + 1}.$$

8.2 *In the Calculus of Limits, the Poisson Kernel does not have*

the sifting property

Proof:

For $\xi - x = 2m$,

$$\begin{aligned} \frac{k-1}{2k(k-1)[1-\cos\pi(x-\xi)]+1} &= \frac{k-1}{2k(k-1)[1-\cos(\pi 2m)]+1} \\ &= k-1 \xrightarrow[k \rightarrow \infty]{} \infty. \square \end{aligned}$$

8.3 Hyper-real Poisson Kernel in Infinitesimal Calculus

$$\mathcal{P}_{oisson}(\xi - x) = \begin{cases} \langle k-1 \rangle, & \xi - x = 2m \\ 0, & \xi - x \neq 2m \end{cases}$$

Proof: At any $\xi - x = 2m$, the Kernel is an infinite hyper-real. \square

At any $\xi - x \neq 2m$,

$$\frac{k-1}{2k(k-1)[1-\cos\pi(\xi-x)]+1} = \frac{1}{2k[1-\cos\pi(\xi-x)]+\frac{1}{k-1}} = \text{infinitesimal}. \square$$

8.4 Let $N = \frac{1}{dr}$ be an infinite Hyper-real, Then

$$\begin{aligned} \mathcal{P}_{oisson}(\xi - x) &= \frac{1}{2} \frac{1-r^2}{1-2r\cos\pi(\xi-x)+r^2} \Big|_{r=1-dr} \\ &= \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos\pi[\xi-x])+(dr)^2} \\ &= \dots + \delta(\xi-x+2) + \delta(\xi-x) + \delta(\xi-x-2) + \dots \\ &= \delta_{periodic}(\xi-x). \end{aligned}$$

Proof:

$$\begin{aligned}
\mathcal{P}_{oisson}(\xi - x) &\equiv \frac{1}{2} \frac{1 - r^2}{1 - 2r \cos \pi(\xi - x) + r^2} \Big|_{r=1-dr} \\
&= \frac{1}{2} \frac{1 - (1 - dr)^2}{1 - 2(1 - dr) \cos \pi[\xi - x] + (1 - dr)^2} \\
&= \frac{1}{2} \frac{dr(2 - dr)}{2(1 - dr)(1 - \cos \pi[\xi - x]) + (dr)^2}
\end{aligned}$$

For $x = 2m$,

$$\begin{aligned}
\mathcal{P}_{oisson}(2m) &= \frac{1}{2} \frac{dr(2 - dr)}{2(1 - dr)(1 - \cos \pi 2m) + (dr)^2} \\
&= \frac{1}{2} \frac{dr(2 - dr)}{(dr)^2} \\
&= \frac{1}{dr} - \frac{1}{2}. \square
\end{aligned}$$

For any $x \neq 2m$,

$$\begin{aligned}
\mathcal{P}_{oisson}(x) &= \frac{1}{2} \frac{dr(2 - dr)}{2(1 - dr)(1 - \cos \pi x) + (dr)^2} \\
&\approx \frac{1}{2} \frac{2dr}{2(1 - \cos \pi x)} \\
&= \text{infinitesimal}. \square
\end{aligned}$$

Therefore,

$$\mathcal{P}_{oisson}(\xi - x) = .. + \left\{ \frac{1}{dr}, \xi - x = -2 \right. + \left\{ \frac{1}{dr}, \xi = x \right. + \left\{ \frac{1}{dr}, \xi - x = 2 \right. + ..$$

$$\begin{aligned}
&= \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots \\
&= \delta_{Periodic}(\xi - x). \square
\end{aligned}$$

8.5

$$\begin{aligned}
&.. + \delta(\xi - x + 4) + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \delta(\xi - x - 4) + .. = \\
&= \frac{1}{2} \frac{dr(2 - dr)}{(1 - dr)(1 - \cos \pi[\xi - x]) + (dr)^2}
\end{aligned}$$

9.

Poisson Integral and $\delta_{Periodic}(\xi - x)$

9.1 Poisson Integral of a Hyper-real Function $f(x)$

Let $f(x)$ be a hyper-real function integrable on $[-1,1]$, so that $f(1) = f(-1)$.

Then, for each $n = \dots, -3, -2, -1, 0, 1, 2, 3, \dots$, the integrals

$$\frac{1}{2} \int_{\xi=-1}^{\xi=1} f(\xi) e^{-in\pi\xi} d\xi \equiv c_n$$

exist, [Dan8], with finite, or infinite hyper-real values. The c_n are the Fourier Coefficients of $f(x)$.

The Fourier Series associated with $f(x)$ is

$$\dots + c_{-n} e^{i(-n)\pi x} + \dots + c_{-1} e^{i(-1)\pi x} + c_0 + c_1 e^{i(1)\pi x} + \dots + c_n e^{i(n)\pi x} + \dots =$$

It may equal The Poisson Integral associated with $f(x)$ at $r = 1 - dr$,

$$\begin{aligned} \mathcal{P}_{oisson} \mathcal{S}\{f(x)\} &\equiv \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1-r^2}{1+r^2-2r\cos\pi(\xi-x)} d\xi \Bigg|_{r=1-dr} \\ &= \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos\pi[\xi-x]) + (dr)^2} d\xi \end{aligned}$$

For each x , it may assume finite or infinite hyper-real values.

$$\mathbf{9.2} \quad \mathcal{P}_{oisson} \mathcal{S} \{ \delta_{Periodic}(\xi - x) \} = \delta_{Periodic}(\xi - x)$$

Proof:

$$\begin{aligned} \mathcal{P}_{oisson} \mathcal{S} \{ \delta_{Periodic}(\xi - x) \} &= \int_{\xi=-1}^{\xi=1} \delta_{Periodic}(\xi) \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi[\xi-x]) + (dr)^2} d\xi \\ &= \int_{\xi=-1}^{\xi=1} \{ \dots + \delta(\xi+2) + \delta(\xi) + \delta(\xi-2) + \dots \} \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi[\xi-x]) + (dr)^2} d\xi \\ &= \dots + \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi[2+x]) + (dr)^2} + \\ &\quad + \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi x) + (dr)^2} + \\ &\quad + \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi[2-x]) + (dr)^2} + \dots \\ &= \dots + \dots + \delta(x+2) + \delta(x) + \delta(x-2) + \dots \\ &= \delta_{Periodic}(x). \square \end{aligned}$$

10.

Poisson Integral Theorem

The Poisson Integral Theorem for a hyper-real function $f(x)$, is the Fundamental Theorem for Abel Summation of Fourier series.

It supplies the conditions under which the Poisson Integral associated with $f(x)$ at $r = 1 - dr$, equals $f(x)$.

It is believed to hold in the Calculus of Limits under some Conditions. In fact,

The Theorem cannot be proved in the Calculus of Limits under any conditions,

because it requires integration of the singular Poisson Kernel.

10.1 Poisson Integral Theorem cannot be proved in the Calculus of Limits

Proof:

In the Calculus of Limits,

$$\lim_{r \uparrow 1} \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(\xi-x)} d\xi = \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \lim_{r \uparrow 1} \underbrace{\frac{1-r^2}{1+r^2-2r \cos \pi(\xi-x)}}_{\text{Poisson Kernel}} d\xi.$$

As $r \uparrow 1$, the Poisson Sequence becomes the Poisson Kernel, which is singular, and diverges at any $\xi - x = 2m$.

$$\xi - x = 2m \Rightarrow \cos \pi(\xi - x) = 1,$$

$$\begin{aligned} \Rightarrow \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \pi(\xi - x)} &= \frac{1}{2} \frac{1 + r}{1 - r}, \\ &\xrightarrow{r \uparrow 1} \frac{1}{2} \frac{1 + 1}{0 +} = \infty \end{aligned}$$

Thus, the Poisson Integral diverges in the Calculus of Limits.

Avoiding the singularity at $\xi = x$, by using the Cauchy Principal Value of the integral does not recover the Theorem, because at any

$$\xi - x \neq 2m,$$

$$\begin{aligned} \frac{1}{2} \frac{1 - r^2}{1 + r^2 - 2r \cos \pi(\xi - x)} &= \frac{1}{2} \frac{1 - r^2}{(1 - r)^2 + 4r \sin^2[\frac{1}{2}\pi(\xi - x)]} \\ &\xrightarrow{r \uparrow 1} \frac{0}{2 \sin^2[\frac{1}{2}\pi(\xi - x)]} = 0 \end{aligned}$$

That is, the Poisson Kernel vanishes, and the integral is identically zero, for any function $f(x)$.

Thus, the Poisson Integral Theorem does not hold in the Calculus of Limits. \square

10.2 Calculus of Limits Conditions are irrelevant to Poisson Integral Theorem

Proof:

The Poisson Conditions are

1. $|f(x)|$ is integrable on $[c - L, c + L]$
2. $f(x)$ is periodic with period $T = 2L$
3. $\frac{1}{2}(f(x + 0) + f(x - 0))$ replaces $f(x)$ at a discontinuity point.

It is clear from 10.1 that the Fejer conditions on $f(x)$ do not resolve the singularity of the Fejer kernel, and are not sufficient for the Fejer Summation Theorem. \square

In Infinitesimal Calculus, by 8.4, the Poisson Kernel is the Periodic Delta Function, and by 9.2, it equals its Poisson Integral at $r = 1 - dr$.

Then, the Poisson Integral Theorem holds for any periodic Hyper-Real Function:

10.3 Poisson Integral Theorem for Hyper-real $f(x)$

If $f(x)$ is hyper-real function integrable on $[c - L, c + L]$, so that

$$f(c - L) = f(c + L)$$

Then,
$$f(x) = \mathcal{P}_{oisson} \mathcal{S} \{ f(x) \}$$

Proof: Take $L = 1$, and $c = 0$.

$$f(x) = \int_{\xi=-1}^{\xi=1} f(\xi) \underbrace{\{ \dots + \delta(\xi - x + 2) + \delta(\xi - x) + \delta(\xi - x - 2) + \dots \}}_{\delta_{Periodic}(\xi-x), \text{ where the period of Delta is } T=2} d\xi .$$

By 8.4, $\delta_{Periodic}(\xi - x) = P_{oisson}(\xi - x)$. Thus,

$$\begin{aligned} f(x) &= \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{dr(2-dr)}{2(1-dr)(1-\cos \pi[\xi-x]) + (dr)^2} d\xi \\ &= \int_{\xi=-1}^{\xi=1} f(\xi) \frac{1}{2} \frac{1-r^2}{1+r^2-2r \cos \pi(\xi-x)} \Big|_{r=1-dr} d\xi \\ &= \mathcal{P}_{oisson} \mathcal{S}\{f(x)\}. \square \end{aligned}$$

In particular, the Periodic Delta Function

violates the Poisson Conditions

- ❖ *The Hyper-real $\delta(x)$, is not defined in the Calculus of Limits, and $|\delta(x)|$ is not integrable in any bounded interval.*
- ❖ *$\frac{1}{2}(\delta(x+0) + \delta(x-0)) = 0$ does not replace $\delta(x)$ at its discontinuity point, $x = 0$.*

But by 9.2, $\delta_{Periodic}(x)$ satisfies the Poisson Integral Theorem.

References

- [Achieser] Achieser, N. I., *Theory of Approximation*, Ungar, 1956.
- [Carslaw] Carslaw, H. S., *“Introduction to the Theory of Fourier Series and integrals”* Third Edition, Macmillan, 1930.
- [Dan1] Dannon, H. Vic, *“Well-Ordering of the Reals, Equality of all Infinities, and the Continuum Hypothesis”* in Gauge Institute Journal Vol. 6 No. 2, May 2010;
- [Dan2] Dannon, H. Vic, *“Infinitesimals”* in Gauge Institute Journal Vol.6 No. 4, November 2010;
- [Dan3] Dannon, H. Vic, *“Infinitesimal Calculus”* in Gauge Institute Journal Vol. 7 No. 4, November 2011;
- [Dan4] Dannon, H. Vic, *“Riemann’s Zeta Function: the Riemann Hypothesis Origin, the Factorization Error, and the Count of the Primes”*, in Gauge Institute Journal of Math and Physics, Vol. 5, No. 4, November 2009.
- [Dan5] Dannon, H. Vic, *“The Delta Function”* in Gauge Institute Journal Vol. 8, No. 1, February, 2012;
- [Dan6] Dannon, H. Vic, *“Delta Function, the Fourier Transform, and the Fourier Integral Theorem”* in Gauge Institute Journal Vol. 8, No. 2, May, 2012;
- [Dan7] Dannon, H. Vic, *“Riemannian Trigonometric Series”*, Gauge Institute Journal, Volume 7, No. 3, August 2011.
- [Dan8] Dannon, H. Vic, *“[Periodic Delta Function and Dirichlet Summation of Fourier Series](#)”* posted to www.gauge-institutue.org;
- [Dan9] Dannon, H. Vic, *“Lebesgue Integration”* Gauge Institute Journal Vol.7 No. 1, February 2011;

[Dan10] Dannon, H. Vic, "[Periodic Delta Function and Fejer-Cesaro Summation of Fourier Series](#)", June 2012, posted to www.gauge-institute.org

[Dan11] Dannon, H. Vic, "*Infinite Series with Infinite Hyper-real Sum*" in Gauge Institute Journal Vol. 8, No. 3, August, 2012;

[Davis] Davis, Philip, "*Interpolation and Approximation*" Blaisdell, 1963.

[Hardy] Hardy, G. H., *Divergent Series*, Chelsea 1991.

[Natanson] Natanson, I. P., "*Constructive Function Theory*" Ungar, 1964.

[Rogosinski] Rogosinski, Werner, "*Fourier Series*" Chelsea, 1950.

[Tolstov] Tolstov, Georgi, "*Fourier Series*" Prentice-Hall, 1962

[Zygmund] Zygmund, A., "*Trigonometric Series*", Second Edition, Cambridge University Press, 1968.